



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra

A local property of basic Rickard equivalences

Lluís Puig^a, Yuanyang Zhou^{b,*}, 1^a CNRS, Université Paris 7, 6 Av Bizet, 94340 Joinville-le-Pont, France^b Department of Mathematics and Statistics, Central China Normal University, Wuhan 430079, PR China

ARTICLE INFO

Article history:

Received 25 July 2008

Available online 17 July 2009

Communicated by Michel Broué

Keywords:

Basic Rickard equivalences

Blocks

ABSTRACT

Let p be a prime and k be an algebraically closed field of characteristic p . Let G and G' be two finite groups, b and b' be blocks of G and G' and (P, e) and (P', e') be maximal Brauer (b, G) - and (b', G') -pairs. If kGb and $kG'b'$ are basic Rickard equivalent, then there is a group isomorphism $\lambda: P \cong P'$ such that it induces an equivalence between the Brauer categories of kGb and $kG'b'$ and that, for any Brauer (b, G) -pair (Q, f) contained in (P, e) and any subgroup K of the image of $N_G(Q, f)$ in $\text{Aut}(Q)$, the block algebras $kN_G^K(Q)f$ and $kN_{G'}^{K'}(Q')f'$ are basic Rickard equivalent, where $Q' = \lambda(Q)$, f' is the block of $kC_{G'}(Q')$ such that $(Q', f') \leq (P', e')$, and K' denotes the image of K in $\text{Aut}(Q')$.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

1.1. Let p be a prime and k an algebraically closed field of characteristic p . If two block algebras kGb and $kG'b'$ of two finite groups G and G' are *basic Morita equivalent* – namely, they are Morita equivalent via an indecomposable $k(G \times G')$ -module having a vertex $\tilde{P} \subset G \times G'$ and a source \tilde{N} such that \tilde{P} stabilizes a k -basis of $\text{End}_k(\tilde{N})$ [6, Section 7] – then there exists a group isomorphism $P \cong P'$ between their defect groups P and P' , inducing an equivalence between their *local categories* [6, 7.6.6]. In this case, we have proved in [7] that for any local pointed group Q_δ on kGb , denoting by $Q'_{\delta'}$ the local pointed group on $kG'b'$ corresponding to Q_δ throughout this equivalence, the respective blocks b_δ and $b_{\delta'}$ of the normalizers $N_G(Q_\delta)$ and $N_{G'}(Q'_{\delta'})$ determined by δ and δ' (see Section 4 below) are *basic Morita equivalent* too, which extends statement 7.7.4 in [6].

* Corresponding author.

E-mail addresses: puig@math.jussieu.fr (L. Puig), zhouy74@163.com (Y. Zhou).

¹ During the preparation of this paper, the second author was supported by NSFC (Grant No. 10501016), the Key Project of Chinese Ministry of Education and the Alexander von Humboldt Foundation of Germany; he thanks a lot for their financial supports.

1.2. Since Jeremy Rickard's thesis [8], it has been understood the interest in considering not only the equivalence between the categories of kGb - and $kG'b'$ -modules, but more generally the equivalence between their corresponding *derived categories*. Thus, in [6, Section 19] are also considered the *basic Rickard equivalences* which extend both, the *splendid equivalences* previously introduced by Rickard [9] and the *basic Morita equivalences* mentioned above. In all these cases, the word “*basic*” means the existence of suitable “*stable bases*” which originally implies the same kind of equivalence at the level of the so-called *Brauer correspondents* in the centralizers [6, Theorem 19.11].

1.3. Our main tool in [7] in order to prove that *basic Morita equivalences* are inherited by the normalizers – more generally, by the K -normalizers – of the corresponding *Brauer correspondents* was the so-called *extended Brauer construction* (see Section 3 below). In this paper, we extend this result to the *basic Rickard equivalences* and, as a matter of fact, our proof only depends on the *extended Brauer construction* again and on the machinery already developed in [6]. A previous result by Andrei Marcus [1] on the same direction only covers the case where K is a p' -group – actually, Marcus' effort amounts to guessing that the equivalence for the centralizers *already forces* the equivalence for the normalizers, which it is not our approach.

1.4. Let us briefly recall the notation from [6] that we need here. In the point of view of [6], the *differential complexes* are considered as \mathfrak{D} -modules where, denoting by \mathfrak{F} the commutative k -algebra of all the k -valued functions on the set \mathbf{Z} of all rational integers, \mathfrak{D} is the k -algebra containing \mathfrak{F} as a unitary k -subalgebra and an element d such that

$$\mathfrak{D} = \mathfrak{F} \oplus \mathfrak{F}d, \quad d^2 = 0 \quad \text{and} \quad df = \text{sh}(f)d \neq 0 \quad \text{for any } f \in \mathfrak{F} - \{0\}$$

where sh denotes the automorphism on the k -algebra \mathfrak{F} mapping $f \in \mathfrak{F}$ onto the k -valued function sending $z \in \mathbf{Z}$ to $f(z+1)$; moreover, we denote by s and i_z , for any $z \in \mathbf{Z}$, the k -valued functions mapping $z' \in \mathbf{Z}$ on $(-1)^{z'}$ and $\delta_z^{z'}$ respectively. Except for all the *group algebras* over \mathfrak{D} , we assume that all the modules and the algebras over k are finite dimensional. If A is a k -algebra we denote by A^* the group of invertible elements of A , and by A° the opposite k -algebra. Note that we have an isomorphism $t: \mathfrak{D} \cong \mathfrak{D}^\circ$ mapping $f \in \mathfrak{F}$ on the k -valued function sending $z \in \mathbf{Z}$ to $f(-z)$, and d on sd .

1.5. A \mathfrak{D} -interior algebra is a k -algebra A endowed with a unitary k -algebra homomorphism $\mathfrak{D} \rightarrow A$. Note that the isomorphism $t: \mathfrak{D} \cong \mathfrak{D}^\circ$ then determines a \mathfrak{D} -interior algebra structure for A° . Moreover, we have a k -algebra homomorphism $\mathfrak{D} \rightarrow k$ mapping $f + f'd$ on $f(0)$ for any $f, f' \in \mathfrak{F}$, so that any k -algebra admits a *trivial* structure of \mathfrak{D} -interior algebra. If A and A' are \mathfrak{D} -interior algebras, the tensor product $A \otimes_k A'$ admits a \mathfrak{D} -interior algebra structure given by

$$\begin{aligned} f \cdot (a \otimes a') &= \sum_{z, z' \in \mathbf{Z}} f(z+z') i_z \cdot a \otimes i_{z'} \cdot a', \\ d \cdot (a \otimes a') &= d \cdot a \otimes s \cdot a' + a \otimes d \cdot a' \end{aligned}$$

which makes sense since in the sum above all but a finite number of terms vanish [6, Proposition 9.7] and since we have $\text{sh}(s) = -s$.

1.6. Let G be a finite group; recall that a kG -interior algebra is a k -algebra endowed with a unitary k -algebra homomorphism from kG . Similarly, a $\mathfrak{D}G$ -interior algebra is a k -algebra A endowed with a unitary k -algebra homomorphism $\rho: \mathfrak{D}G \rightarrow A$ (but A is always finite dimensional!); for any $x \in \mathfrak{D}G$ and $a \in A$, we write $x \cdot a$ and $a \cdot x$ instead of $\rho(x)a$ and $a\rho(x)$ respectively. For any subgroup H of G , we denote by A^H the centralizer of $\rho(H)$ in A ; obviously $\rho(x) \in A^H$ for any $x \in \mathfrak{D}C_G(H)$ and thus the restriction of ρ to $\mathfrak{D}C_G(H)$ induces a $\mathfrak{D}C_G(H)$ -interior algebra structure on A^H . If $\varphi: L \rightarrow G$ is a group homomorphism, the composition of the corresponding k -algebra homomorphism $\mathfrak{D}L \rightarrow \mathfrak{D}G$

with $\rho: \mathfrak{D}G \rightarrow A$ defines a $\mathfrak{D}L$ -interior algebra structure on A ; we denote this $\mathfrak{D}L$ -interior algebra by $\text{Res}_\varphi(A)$; we write $\text{Res}_H^G(A)$ when $L = H$ and φ is the inclusion homomorphism $H \subset G$.

1.7. Let us denote by $\mathbb{C}_0(A)$ the centralizer of the image of \mathfrak{D} in A ; since the images of \mathfrak{D} and G centralize each other, $\mathbb{C}_0(A)$ inherits a kG -interior algebra structure and, according to the terminology in [6], the *pointed groups, their inclusions, the local pointed groups, etc.*, over the $\mathfrak{D}G$ -interior algebra A are nothing but the pointed groups, their inclusions, the local pointed groups, etc., over the kG -interior algebra $\mathbb{C}_0(A)$. However, if H_β is a pointed group over A , so that β is a conjugacy class of primitive idempotents in $\mathbb{C}_0(A)^H$, for any $i \in \beta$ the k -algebra $A_\beta = iAi$ inherits a $\mathfrak{D}H$ -interior algebra structure mapping $y \in \mathfrak{D}H$ on $y \cdot i = i \cdot y$. For any subgroup H of G , we call *contractible* any point contained in the two-sided ideal

$$\mathbb{B}_0(A^H) = \mathbb{C}_0(A)^H \cap \{d \cdot a + a \cdot d \mid a \in A^H\}$$

and we set $\mathbb{H}_0(A^H) = \mathbb{C}_0(A)^H / \mathbb{B}_0(A^H)$, which still inherits a $kC_G(H)$ -interior algebra structure; whenever $\mathbb{H}_0(A^G) = \{0\}$ we say that A is *contractible*. It is clear that if M is a $\mathfrak{D}G$ -module then $\text{End}_k(M)$ is a $\mathfrak{D}G$ -interior algebra and we say that M is *contractible* whenever $\text{End}_k(M)$ is so [6, Corollary 10.9]; moreover, we say that M is *0-split* if it is $\mathfrak{D}G$ -isomorphic to the direct sum of a contractible $\mathfrak{D}G$ -module and a kG -module endowed with the trivial \mathfrak{D} -structure defined above [6, 10.12].

1.8. Our standard setting in this paper is formed by two finite groups G and G' , respective blocks b of G and b' of G' , and an indecomposable $\mathfrak{D}(G \times G')$ -module \check{M} associated with $b \otimes b'$ such that the restrictions of \check{M} to $G \times \{1\}$ and to $\{1\} \times G'$ are both projective. We denote by \check{M}^* the k -dual of \check{M} which, via the isomorphism t (cf. 1.4), still has a $\mathfrak{D}(G \times G')$ -module structure. Following [6, 18.2.2], we say that \check{M} defines a *Rickard equivalence* between kGb and $kG'b'$ if, for suitable contractible $\mathfrak{D}(G \times G')$ - and $\mathfrak{D}(G' \times G)$ -modules C and C' , we have respective $\mathfrak{D}(G \times G)$ - and $\mathfrak{D}(G' \times G')$ -module isomorphisms

$$\check{M} \otimes_{kG'} \check{M}^* \cong kGb \oplus C \quad \text{and} \quad \check{M}^* \otimes_{kG} \check{M} \cong kG'b' \oplus C'$$

where kGb and $kG'b'$ have the trivial \mathfrak{D} -interior structure defined above. We say that kGb and $kG'b'$ are *Rickard equivalent* if there exists such a $\mathfrak{D}(G \times G')$ -module; in this case, note that the $\mathfrak{D}(G \times G)$ - and $\mathfrak{D}(G' \times G')$ -modules $\check{M} \otimes_{kG'} \check{M}^*$ and $\check{M}^* \otimes_{kG} \check{M}$ are 0-split.

1.9. By our remarks in 1.7 above, in this case we still have a vertex \check{P} of \check{M} — we consider a maximal *local pointed group* $\check{P}_{\check{y}}$ over the $\mathfrak{D}(G \times G')$ -interior algebra $\text{End}_k(\check{M})$ or, equivalently, over the $k(G \times G')$ -interior algebra $\mathbb{C}_0(\text{End}_k(\check{M}))$ — and a corresponding *source* \check{N} — the $\mathfrak{D}\check{P}$ -module $j \cdot \check{M}$ for some $j \in \check{y}$. According to Theorem 18.8 in [6], the images $P \subset G$ and $P' \subset G'$ of $\check{P} \subset G \times G'$ by the canonical projections $\pi: G \times G' \rightarrow G$ and $\pi': G \times G' \rightarrow G'$ are *defect groups* of b and b' respectively; then, it is clear that $\check{P} \times \check{P}$ acts on $\text{End}_k(\check{N})$ by left and right multiplication, and we say that the Rickard equivalence between kGb and $kG'b'$ defined by \check{M} is *basic* whenever each one of the subgroups $\check{P} \times_P \check{P}$ and $\check{P} \times_{P'} \check{P}$ of $\check{P} \times \check{P}$ stabilizes a basis of $\text{End}_k(\check{N})$. We say that kGb and $kG'b'$ are *basic Rickard equivalent* if there exists a $\mathfrak{D}(G \times G')$ -module \check{M} defining a *basic Rickard equivalence* between kGb and $kG'b'$.

1.10. In this situation, as in the case of the *basic Morita equivalences*, we have a group isomorphism $\lambda: P \cong P'$ but this time λ only induces an equivalence between the *Brauer categories* of kGb and $kG'b'$ [6, Theorem 19.7]. That is to say, recall that a *Brauer* (b, G) -pair (Q, f) is a pair formed by a subgroup Q of G and a block f of $C_G(Q)$ such that $f \text{Br}_Q(b) = f$ (cf. 3.1 below); since $(kG)(Q) \cong kC_G(Q)$ (cf. 3.1 below), any local point δ of Q over kGb determines a Brauer (b, G) -pair (Q, f) and it is well-known that the inclusion between the local pointed groups over kGb induces an inclusion relation between the Brauer (b, G) -pairs [6, 2.13]; then, the *Brauer category* of kGb is formed by the Brauer (b, G) -pairs and by the homomorphisms between the groups induced by the

G -conjugation and the inclusion of (b, G) -Brauer pairs. Recall that (Q, f) is called *selfcentralizing* if Q is a defect group of f as a block of $Q \cdot C_G(Q)$ [6, 2.12]; then, Q has a unique local point δ over kGb associated with f [6, 2.12.1] and Q_δ is called *selfcentralizing* too.

1.11. Moreover recall that, if (P, e) and (P', e') are Brauer (b, G) - and (b', G') -pairs, for any subgroup Q of P and any subgroup Q' of P' there are unique Brauer (b, G) - and (b', G') -pairs fulfilling [6, 2.13.2]

$$(Q, f) \subset (P, e) \quad \text{and} \quad (Q', f') \subset (P', e')$$

and that the corresponding *full subcategories* over all these objects are equivalent to the respective *Brauer categories* of kGb and $kG'b'$. Then, Theorem 19.7 in [6] states that, if kGb and $kG'b'$ are *basic Rickard equivalent*, there is a group isomorphism $\lambda: P \cong P'$ such that the correspondence mapping any Brauer (b, G) -pair $(Q, f) \subset (P, e)$ on the unique Brauer (b', G') -pair $(\lambda(Q), f^\lambda) \subset (P', e')$ induces an equivalence between the *Brauer categories* of kGb and $kG'b'$. Finally, if Q is a subgroup of P and K a subgroup of $\text{Aut}(Q)$, let us denote by $N_G^K(Q)$ the converse image of K in $N_G(Q)$ – called *K-normalizer* of Q in G . We are ready to state our main result.

Theorem 1.12. Assume that kGb and $kG'b'$ are basic Rickard equivalent. Let (P, e) and (P', e') be respective maximal Brauer (b, G) - and (b', G') -pairs. Then, there is a group isomorphism $\lambda: P \cong P'$ such that it induces an equivalence between the Brauer categories of kGb and $kG'b'$ and that, for any Brauer (b, G) -pair (Q, f) contained in (P, e) and any subgroup K of the image of $N_G(Q, f)$ in $\text{Aut}(Q)$, the block algebras $kN_G^K(Q)f$ and $kN_G^{K'}(Q')f'$ are basic Rickard equivalent, where $Q' = \lambda(Q)$, $f' = f^\lambda$, and K' denotes the image of K in $\text{Aut}(Q')$.

2. Notation and quoted results

2.1. Let G be a finite group and A a $\mathcal{D}G$ -interior algebra; as we mention in 1.7 above, a *pointed group* H_β over A is nothing but a pointed group over the kG -interior algebra $\mathbb{C}_0(A)$, namely β is a conjugacy class of primitive idempotents in $\mathbb{C}_0(A)^H = \mathbb{C}_0(A^H)$; similarly, a pointed group P_γ over A is contained in H_β if $P \subset H$ and for any $i \in \beta$ there is $j \in \gamma$ such that $ij = j = ji$; we say that the point γ is *local* and that P_γ is a *local pointed group* if $\gamma \not\subset \sum_Q \mathbb{C}_0(A)_Q^P$ when Q runs over the set of proper subgroups of P .

2.2. A *homomorphism* $f: A \rightarrow A'$ between $\mathcal{D}G$ -interior algebras is a not necessarily unitary k -algebra homomorphism fulfilling $f(x \cdot a \cdot y) = x \cdot f(a) \cdot y$ for any $x, y \in \mathcal{D}$ and any $a \in A$; then, we say that f is an *embedding* whenever $\text{Ker}(f) = \{0\}$ and $\text{Im}(f) = f(1) \cdot A' \cdot f(1)$. Note that any element $a \in \mathbb{C}_0(A^G)^*$ induces by conjugation a $\mathcal{D}G$ -interior algebra automorphism $\text{int}(a): A \cong A$ mapping $c \in A$ on aca^{-1} ; we set

$$\tilde{f} = \{\text{int}(a') \circ f \mid a' \in \mathbb{C}_0((A')^G)^*\}$$

and call it the *exomorphism* of $\mathcal{D}G$ -interior algebras determined by f . For instance, for any pointed group H_β over A and any $i \in \beta$, the inclusion $iAi \subset A$ determines a canonical exoembedding $\tilde{f}_\beta: A_\beta \rightarrow \text{Res}_H^G(A)$ which, up to a unique $\mathcal{D}H$ -interior algebra exoisomorphism, does not depend on the choice of i .

2.3. Let H be a subgroup of G and B a $\mathcal{D}H$ -interior algebra; the *induced algebra*

$$\text{Ind}_H^G(B) = kG \otimes_{kH} B \otimes_{kH} kG$$

where, for any $x, y, x', y' \in H$ and any $b, b' \in B$, the product is defined by

$$(x \otimes b \otimes y)(x' \otimes b' \otimes y') = \begin{cases} x \otimes b \cdot yx' \cdot b' \otimes y' & \text{if } yx' \in H, \\ 0 & \text{otherwise} \end{cases}$$

clearly admits a $\mathfrak{D}G$ -interior structure mapping $z \in \mathfrak{D}G$ onto $\sum_y zy \otimes 1 \otimes y^{-1}$ where $y \in G$ runs over a set of representatives for G/H ; let us denote by

$$d_H^G(B) : B \rightarrow \text{Res}_H^G(\text{Ind}_H^G(B)) \quad (2.3.1)$$

the $\mathfrak{D}H$ -interior algebra embedding mapping $b \in B$ onto $1 \otimes b \otimes 1$. In order to formulate the main feature of the induced algebras, it is handy to replace A by its *Higman envelope* [6, 14.6]; namely, it follows from Proposition 14.7 and Corollary 14.11 in [6] that we may assume that

2.3.2 For any pointed group H_β over A there is a unique $\mathfrak{D}G$ -interior algebra exoembedding

$$\tilde{h}_\beta : \text{Ind}_H^G(A_\beta) \rightarrow \text{Res}_H^G(A)$$

$$\text{fulfilling } \text{Res}_H^G(\tilde{h}_\beta) \circ \tilde{d}_H^G(A_\beta) = \tilde{f}_\beta.$$

Then, Theorem 14.9 in [6] states.

Theorem 2.4. For any pointed groups H_β and L_ε over A fulfilling $L \subset H$, the following two conditions are equivalent:

2.4.1 $\beta \subset \text{Tr}_L^H(\mathbb{C}_0(A^L) \cdot \varepsilon \cdot \mathbb{C}_0(A^L))$.

2.4.2 There is a $\mathfrak{D}H$ -exoembedding $\tilde{h}_\beta^\varepsilon : A_\beta \rightarrow \text{Ind}_L^H(A_\varepsilon)$ fulfilling

$$\tilde{h}_\varepsilon \circ \text{Ind}_H^G(\tilde{h}_\beta^\varepsilon) = \tilde{h}_\beta.$$

Moreover, in this case the $\mathfrak{D}H$ -exoembedding $\tilde{h}_\beta^\varepsilon$ is unique.

2.5. But, we also need the *noninjective induction* introduced in [6]; let us recall its definition. Let \bar{G} be another finite group and $\rho : G \rightarrow \bar{G}$ be a surjective group homomorphism with a kernel W ; the tensor product $k \otimes_{k_W} A$ obviously admits a right A -module structure

$$(k \otimes_{k_W} A) \times A \rightarrow k \otimes_{k_W} A.$$

Since we have the equality $(1 \otimes a) \cdot b = (1 \otimes a) \cdot (x \cdot b)$ for any $a \in A$ such that $1 \otimes a \in (k \otimes_{k_W} A)^W$, any $b \in A$ and any $x \in W$, the restriction to $(k \otimes_{k_W} A)^W \times A$ of the above map factorizes throughout the maps

$$\begin{aligned} (k \otimes_{k_W} A)^W \times A &\rightarrow (k \otimes_{k_W} A)^W \times (k \otimes_{k_W} A), \\ (k \otimes_{k_W} A)^W \times (k \otimes_{k_W} A) &\rightarrow k \otimes_{k_W} A \end{aligned}$$

and therefore the restriction of the latter to $(k \otimes_{k_W} A)^W \times (k \otimes_{k_W} A)^W$ induces a product on $(k \otimes_{k_W} A)^W$

$$(k \otimes_{k_W} A)^W \times (k \otimes_{k_W} A)^W \rightarrow (k \otimes_{k_W} A)^W.$$

With this product and the homomorphism $\mathfrak{D}\bar{G} \rightarrow (k \otimes_{k_W} A)^W$ sending $z \in \mathfrak{D}$ to $1 \otimes (z \cdot 1_A)$ and $\bar{x} \in \bar{G}$ to $1 \otimes (x \cdot 1_A)$ where x is a lifting of \bar{x} to G , $(k \otimes_{k_W} A)^W$ becomes a $\mathfrak{D}\bar{G}$ -interior algebra [6, 3.2.5] – noted $\text{Ind}_\varphi(A)$.

2.6. A significant difference from the ordinary induction is that we have no canonical homomorphism from A , but only from the *normalizer* of W in A . Precisely, for any $\sigma \in \text{Aut}(W)$, let us denote by $\Delta_\sigma : W \rightarrow W \times W$ the σ -twisted diagonal homomorphism mapping $w \in W$ on $(\sigma(w), w)$ and set

$$N_A(W) = \bigoplus_{\sigma \in \text{Aut}(W)} A^{\Delta_\sigma(W)}$$

which admits a $\mathfrak{D}G$ -interior algebra structure (see [7, Section 3] and Section 3 below); moreover, we have an evident unitary $\mathfrak{D}G$ -interior algebra homomorphism $N_A(W) \rightarrow A$ which, as it is not difficult to prove, is injective whenever A is projective as $k(W \times W)$ -module (and then $N_A(W)$ coincides with the definition in [6, 2.3.3]). Then, we denote by [6, 3.4.2]

$$d_\rho(A) : N_A(W) \rightarrow \text{Res}_\rho(\text{Ind}_\rho(A))$$

the homomorphism mapping $a \in A^{\Delta_\sigma(W)}$ on $1 \otimes a$ for any $\sigma \in \text{Aut}(W)$. More generally, for any group homomorphism $\varphi : G \rightarrow G'$, setting $\bar{G} = \varphi(G)$ and denoting by $\rho : G \rightarrow \bar{G}$ the group homomorphism determined by φ , we still set

$$\text{Ind}_\varphi(A) = \text{Ind}_G^{G'}(\text{Ind}_\rho(A)) \quad \text{and} \quad d_\varphi(A) = d_G^{G'}(\text{Ind}_\rho(A)) \circ d_\rho(A).$$

2.7. Let us come back to our standard setting (cf. 1.8) and respectively denote by $\pi_{\tilde{P}} : \tilde{P} \rightarrow G$ and $\pi'_{\tilde{P}} : \tilde{P} \rightarrow G'$ the restrictions to \tilde{P} of π and π' . As in [6, 16.1], it follows from Proposition 14.7 and Corollary 14.12 in [6] that we may replace \tilde{N} by a bigger $\mathfrak{D}\tilde{P}$ -module \tilde{N}' — with its restrictions to $\text{Ker}(\pi_{\tilde{P}})$ and $\text{Ker}(\pi'_{\tilde{P}})$ always projective, and containing \tilde{N} as a direct summand — in such a way that $\tilde{S} = \text{End}_k(\tilde{N}')$ becomes a $\mathfrak{D}\tilde{P}$ -interior algebra which coincides with its *Higman envelope* (cf. 2.3.2), and admits $\tilde{P} \times_P \tilde{P}$ - and $\tilde{P} \times_{P'} \tilde{P}$ -stable bases whenever $\text{End}_k(\tilde{N})$ does; then, setting

$$\tilde{A} = \text{Ind}_{\tilde{P}}^{G \times G'}(\tilde{S}), \quad \hat{A} = (\tilde{A})^{1 \times G'} \quad \text{and} \quad \hat{A}' = (\tilde{A})^{G \times 1}$$

we already know that all these interior algebras coincide with their *Higman envelopes* [6, 16.1.3] and that we have $\mathfrak{D}G$ - and $\mathfrak{D}G'$ -interior algebra isomorphisms [6, 16.1.2]

$$\hat{A} \cong \text{Ind}_{\pi_{\tilde{P}}}(\tilde{S} \otimes_k \text{Res}_{\pi'_{\tilde{P}}}(kG')) \quad \text{and} \quad \hat{A}' \cong \text{Ind}_{\pi'_{\tilde{P}}}(\tilde{S} \otimes_k \text{Res}_{\pi_{\tilde{P}}}(kG)).$$

Then, Proposition 18.4 in [6] states.

Proposition 2.8. *With the notation above, \tilde{M} defines a Rickard equivalence between kGb and $kG'b'$ if and only if there are points $\hat{\alpha}$ of G over \hat{A} and $\hat{\alpha}'$ of G' over \hat{A}' such that $\hat{A}_{\hat{\alpha}}$ and $\hat{A}'_{\hat{\alpha}'}$, respectively considered as $\mathfrak{D}(G \times G)$ - and $\mathfrak{D}(G' \times G')$ -modules are 0-split and the structural homomorphisms $kGb \rightarrow \hat{A}_{\hat{\alpha}}$ and $kG'b' \rightarrow \hat{A}'_{\hat{\alpha}'}$ induce kG - and kG' -interior algebra isomorphisms*

$$kGb \cong \mathbb{H}_0(\hat{A}_{\hat{\alpha}}) \quad \text{and} \quad kG'b' \cong \mathbb{H}_0(\hat{A}'_{\hat{\alpha}'}).$$

In this case, the structural homomorphisms induce bijections between the set of pointed groups over kGb and $kG'b'$, and the corresponding sets of noncontractible points over $\hat{A}_{\hat{\alpha}}$ and $\hat{A}'_{\hat{\alpha}'}$, preserving inclusion and localness.

2.9. In [6] the main tool to relate both sides in a Rickard equivalence are the so-called *local tracing triples*. In the above setting, a *local tracing triple* over \hat{A} , \hat{A} and $kG'b'$ is a triple of local pointed groups $Q_{\hat{\delta}}$ on \hat{A} , $\tilde{Q}_{\tilde{\delta}}$ and $Q'_{\delta'}$ on $kG'b'$, such that $\pi(\tilde{Q}) = Q$ and $\pi'(\tilde{Q}) = Q'$, and that, denoting by $\tau: \tilde{Q} \rightarrow Q$ and $\tau': \tilde{Q} \rightarrow Q'$ the corresponding group homomorphisms, there is a (unique) $\mathfrak{D}Q$ -interior algebra exoembedding

$$\tilde{h}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}: \hat{A}_{\hat{\delta}} \rightarrow \text{Ind}_{\tau}(\tilde{A}_{\tilde{\delta}} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}) \quad (2.9.1)$$

in such a way that the following diagram is commutative [6, 16.5.2]

$$\begin{array}{ccc} \text{Res}_{Q \times_{G'}}^{G \times G'}(\tilde{A}) & \xleftarrow{\tilde{h}_{\tilde{\delta}}} & \text{Ind}_{\tilde{Q}}^{Q \times G'}(\tilde{A}_{\tilde{\delta}}) \\ \cup & & \cup \\ \text{Res}_Q^G(\hat{A}) & & \text{Ind}_{\tilde{Q}}^{Q \times G'}(\tilde{A}_{\tilde{\delta}})^{1 \times G'} \cong \text{Ind}_{\tau}(\tilde{A}_{\tilde{\delta}} \otimes_k \text{Res}_{\pi'_Q}(kG')) \\ \tilde{f}_{\tilde{\delta}} \uparrow & & \uparrow \text{Ind}_{\tau}(\text{id} \otimes_k \text{Res}_{\tau'}(\tilde{f}_{\delta'})) \\ \hat{A}_{\hat{\delta}} & \xrightarrow{\tilde{h}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}} & \text{Ind}_{\tau}(\tilde{A}_{\tilde{\delta}} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}) \end{array}$$

explicitly, $\tilde{h}_{\tilde{\delta}}$ induces an exoembedding from $\text{Ind}_{\tilde{Q}}^{Q \times G'}(\tilde{A}_{\tilde{\delta}})^{1 \times G'}$ to $\text{Res}_Q^G(\hat{A})$ and therefore, by composition, we obtain an exoembedding

$$\tilde{g}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}: \text{Ind}_{\tau}(\tilde{A}_{\tilde{\delta}} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}) \rightarrow \text{Res}_Q^G(\hat{A})$$

so that $\tilde{h}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'} \circ \tilde{g}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'} = \tilde{f}_{\tilde{\delta}}$; thus, the existence of $\tilde{h}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}$ is equivalent to the equality

$$f_{\hat{\delta}}(1)g_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}(1) = f_{\hat{\delta}}(1) = g_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}(1)f_{\hat{\delta}}(1) \quad (2.9.2)$$

for a suitable choice of representatives $f_{\hat{\delta}}$ and $g_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}$.

2.10. Moreover, we consider an *inclusion relation* between the local tracing triples refining the corresponding inclusions between the respective local pointed groups; we say that a local tracing triple $(R_{\hat{\varepsilon}}, \tilde{R}_{\tilde{\varepsilon}}, R'_{\varepsilon'})$ is *contained in* $(Q_{\hat{\delta}}, \tilde{Q}_{\tilde{\delta}}, Q'_{\delta'})$ and we write

$$(R_{\hat{\varepsilon}}, \tilde{R}_{\tilde{\varepsilon}}, R'_{\varepsilon'}) \subset (Q_{\hat{\delta}}, \tilde{Q}_{\tilde{\delta}}, Q'_{\delta'})$$

if we have $R_{\hat{\varepsilon}} \subset Q_{\hat{\delta}}$, $\tilde{R}_{\tilde{\varepsilon}} \subset \tilde{Q}_{\tilde{\delta}}$ and $R'_{\varepsilon'} \subset Q'_{\delta'}$ and there is a $\mathfrak{D}R$ -interior algebra exoembedding

$$\tilde{g}_{\tilde{\varepsilon}, \varepsilon'}^{\tilde{\delta}, \delta'}: \text{Ind}_{\rho}(\tilde{A}_{\tilde{\varepsilon}} \otimes_k \text{Res}_{\rho'}(kG')_{\varepsilon'}) \rightarrow \text{Res}_R^Q(\text{Ind}_{\tau}(\tilde{A}_{\tilde{\delta}} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}))$$

fulfilling

$$\tilde{g}_{\tilde{\varepsilon}, \varepsilon'}^{\tilde{\delta}, \delta'} = \text{Res}_R^Q(\tilde{g}_{\tilde{\delta}, \delta'}^{\tilde{\delta}, \delta'}) \circ \tilde{g}_{\tilde{\varepsilon}, \varepsilon'}^{\tilde{\delta}, \delta'}$$

where $\rho: \tilde{R} \rightarrow R$ and $\rho': \tilde{R} \rightarrow R'$ are the respective restrictions of τ and τ' . Then, Theorem 16.15 in [6] guarantees that, for any local pointed group R_{ε} contained in Q_{δ} there is a local tracing triple $(R_{\hat{\varepsilon}}, \tilde{R}_{\tilde{\varepsilon}}, R'_{\varepsilon'})$ contained in $(Q_{\hat{\delta}}, \tilde{Q}_{\tilde{\delta}}, Q'_{\delta'})$.

2.11. Let us come to the “basic” situations. We say that a local tracing triple $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ over \hat{A} , \ddot{A} and $kG'b'$ is *basic* [6, 17.1] whenever \ddot{A}_{δ} is a projective $k(\text{Ker}(\tau) \times \text{Ker}(\tau))$ -module, the subgroup $\ddot{Q} \times_Q \ddot{Q}$ of $\ddot{Q} \times \ddot{Q}$ stabilizes a basis of \ddot{A}_{δ} and \hat{A}_{δ} considered as a $\mathfrak{D}(Q \times Q)$ -module is 0-split (actually, the last condition is stronger than in [6, 17.1]). In this case, τ admits a section and, for any section $\mu: Q \rightarrow \ddot{Q}$ of τ , let us set $Q^{\mu} = \mu(Q)$ and $Q^{\mu'} = \tau(Q^{\mu})$, and let us denote by

$$Z(Q) \xleftarrow{\tau_{\mu}} C_{\ddot{Q}}(Q^{\mu}) \xrightarrow{\tau_{\mu'}} Z(Q^{\mu'})$$

the respective restrictions of τ and τ' ; then, it is clear that the $\mathfrak{D}\ddot{Q}$ -interior algebra $\ddot{A}_{\delta} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}$ is also a projective $k(\text{Ker}(\tau) \times \text{Ker}(\tau))$ -module and that the subgroup $\ddot{Q} \times_Q \ddot{Q}$ of $\ddot{Q} \times \ddot{Q}$ stabilizes a basis on it; hence, according to Theorem 13.9 in [6] applied to this $\mathfrak{D}\ddot{Q}$ -interior algebra and to Q , for any section μ of τ we have a $\mathfrak{D}Z(Q)$ -interior algebra embedding

$$\begin{array}{c} \text{Ind}_{\tau}(\ddot{A}_{\delta} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'})(Q) \\ \uparrow e_{\mu}^{\ddot{\delta}, \delta'} \\ \text{Ind}_{\tau_{\mu}}(\ddot{A}_{\delta}(Q^{\mu}) \otimes_k \text{Res}_{\tau'_{\mu}}((kG')_{\delta'}(Q^{\mu'}))) \end{array}$$

and the set of idempotents $e_{\mu}^{\ddot{\delta}, \delta'}(1)$ form an orthogonal decomposition of the unity in the top k -algebra. We say that a section μ of τ is a *section* of the basic local tracing triple $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ if we have a factorization [6, 17.4]

$$\begin{array}{ccc} \hat{A}_{\delta}(Q) & \xrightarrow{\tilde{h}_{\delta}^{\ddot{\delta}, \delta'}(Q)} & \text{Ind}_{\tau}(\ddot{A}_{\delta} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'})(Q) \\ & \searrow \tilde{h}_{\delta}^{\mu} & \uparrow \tilde{e}_{\mu}^{\ddot{\delta}, \delta'} \\ & \text{Ind}_{\tau_{\mu}}(\ddot{A}_{\delta}(Q^{\mu}) \otimes_k \text{Res}_{\tau'_{\mu}}((kG')_{\delta'}(Q^{\mu'}))) & \end{array}$$

where we set $Q^{\mu} = \mu(Q)$ and $Q^{\mu'} = \tau'(Q^{\mu})$, and denote by τ_{μ} and τ'_{μ} the respective restrictions of τ and τ' .

2.12. In this case, \tilde{h}_{δ}^{μ} is a unique exoembedding [6, 2.11.4] and it follows from the equality $\tilde{g}_{\delta, \delta'}^{\ddot{\delta}} \circ \tilde{h}_{\delta}^{\ddot{\delta}, \delta'} = \tilde{f}_{\delta}$ that

$$\tilde{g}_{\delta, \delta'}^{\ddot{\delta}}(Q) \circ \tilde{e}_{\mu}^{\ddot{\delta}, \delta'} \circ \tilde{h}_{\delta}^{\mu} = \tilde{g}_{\delta, \delta'}^{\ddot{\delta}}(Q) \circ \tilde{h}_{\delta}^{\ddot{\delta}, \delta'}(Q) = \tilde{f}_{\delta}(Q).$$

Conversely, assume that Q_{δ} , \ddot{Q}_{δ} and $Q'_{\delta'}$ are respective local pointed groups on \hat{A} , \ddot{A} and kG' such that \ddot{A}_{δ} is a projective $k(\text{Ker}(\tau) \times \text{Ker}(\tau))$ -module, that the subgroup $\ddot{Q} \times_Q \ddot{Q}$ of $\ddot{Q} \times \ddot{Q}$ stabilizes a basis of \ddot{A}_{δ} , that \hat{A}_{δ} is 0-split considered as a $\mathfrak{D}(Q \times Q)$ -module and that, for some section μ of τ , there is a $\mathfrak{D}Z(Q)$ -interior algebra exoembedding

$$\tilde{h}: \hat{A}_{\delta}(Q) \rightarrow \text{Ind}_{\tau_{\mu}}(\ddot{A}_{\delta}(Q^{\mu}) \otimes_k \text{Res}_{\tau'_{\mu}}((kG')_{\delta'}(Q^{\mu'})))$$

fulfilling

$$\tilde{g}_{\delta, \delta'}(Q) \circ \tilde{e}_{\mu}^{\delta, \delta'} \circ \tilde{h} = \tilde{f}_{\delta}(Q);$$

then $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ is a basic local tracing triple, since the equality (2.9.2) can be easily obtained from this one. Moreover, it follows from Corollary 17.9 in [6] that

2.12.1 if μ and μ' are sections of $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$, there is $x' \in G'$ such that $(1, x')$ normalizes \ddot{Q} and that, denoting by $\varphi_{(1, x')}$ the action of $(1, x')$ on \ddot{Q} , we have $\mu' = \varphi_{(1, x')} \circ \mu$.

2.13. For any subgroup R of Q , it is clear that the restriction of a section μ of $\tau: \ddot{Q} \rightarrow Q$ to a subgroup \tilde{R} of $\tau^{-1}(R)$ such that $\mu(R) \subset \tilde{R}$ is also a section of the restriction $\rho: \tilde{R} \rightarrow R$ of τ ; obviously, $\tilde{A}_{\tilde{R}}$ is still a projective $k(\text{Ker}(\rho) \times \text{Ker}(\rho))$ -module and the subgroup $\tilde{R} \times_R \tilde{R}$ of $\tilde{R} \times \tilde{R}$ stabilizes a basis of $\tilde{A}_{\tilde{R}}$; similarly, if R has a local point \hat{e} on \hat{A} such that $R_{\hat{e}} \subset Q_{\delta}$, $\hat{A}_{\hat{e}}$ considered as a $\mathfrak{D}(R \times R)$ -module is 0-split. In particular, a local tracing triple $(R_{\hat{e}}, \tilde{R}_{\hat{e}}, R'_{\hat{e}'})$ on \hat{A} , \tilde{A} and kG' contained in $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ is basic too and we are interested in choosing it in such a way that the restriction ν to R of a section μ of $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ remains a section of $(R_{\hat{e}}, \tilde{R}_{\hat{e}}, R'_{\hat{e}'})$.

Theorem 2.14. Let $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ be a basic local tracing triple on \hat{A} , \tilde{A} and kG' . For any local pointed group $R_{\hat{e}}$ on \hat{A} contained in Q_{δ} , there is a basic local tracing triple $(R_{\hat{e}}, \tilde{R}_{\hat{e}}, R'_{\hat{e}'})$ contained in $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$ such that the restriction ν of μ to R remains a section of $(R_{\hat{e}}, \tilde{R}_{\hat{e}}, R'_{\hat{e}'})$.

Proof. Arguing by induction on $|Q : R|$, we may assume that $R_{\hat{e}}$ is normal in Q_{δ} . Let μ be a section of $(Q_{\delta}, \ddot{Q}_{\delta}, Q'_{\delta'})$, set

$$\ddot{C}_{\delta, \delta'} = \ddot{A}_{\delta} \otimes_k \text{Res}_{\tau'}(kG')_{\delta'}, \quad \ddot{T} = \tau^{-1}(R) \quad \text{and} \quad T' = \tau'(\ddot{T}),$$

and denote by $\theta: \ddot{T} \rightarrow R$ and $\theta': \ddot{T} \rightarrow T'$ the respective restrictions of τ and τ' , by $\nu: R \rightarrow \ddot{T}$ the restriction of μ , by R^{ν} the image of R in \ddot{T} and by τ_{ν} the corresponding restriction of τ ; it follows from Proposition 13.16 in [6] that we have the commutative diagram of \mathfrak{D} -algebra exoembeddings

$$\begin{array}{ccc} \hat{A}_{\delta}(Q) & \xrightarrow{\tilde{h}_{\delta}^{\delta, \delta'}(Q)} & (\text{Ind}_{\tau}(\ddot{C}_{\delta, \delta'}))(Q) = (\text{Ind}_{\tau'}(\ddot{C}_{\delta, \delta'}))(Q) \\ & \searrow \tilde{h}_{\delta}^{\mu} & \nearrow \tilde{e}_{\mu}^{\delta, \delta'} \\ & \text{Ind}_{\tau_{\mu}}(\ddot{C}_{\delta, \delta'}(Q^{\mu})) & \\ & \searrow & \nearrow \\ & (\text{Ind}_{\tau_{\nu}}(\ddot{C}_{\delta, \delta'}(R^{\nu}))) & (Q/R) \end{array}$$

Then, Theorem 13.9 in [6] applied to $\ddot{C}_{\delta, \delta'}$ with respect to R and the corresponding Brauer homomorphisms [6, Lemma 7.10] determines the following commutative diagram

$$\begin{array}{ccc}
 \hat{A}_{\delta}(R)^Q & \xrightarrow{\quad} & (\text{Ind}_{\tau}(\ddot{C}_{\delta,\delta'}))(R)^Q \\
 \searrow & & \nearrow \\
 \cap & \text{Ind}_{\tau_{\nu}}(\ddot{C}_{\delta,\delta'}(R^{\nu}))^Q & \cap \\
 & \tilde{h}_{\delta}^{\delta,\delta'}(R) & \\
 \hat{A}_{\delta}(R) & \xrightarrow{\quad} & (\text{Ind}_{\tau}(\ddot{C}_{\delta,\delta'}))(R) \\
 \searrow & & \nearrow \\
 & \cap & \\
 & \text{Ind}_{\tau_{\nu}}(\ddot{C}_{\delta,\delta'}(R^{\nu})) & \tilde{e}_{\nu}^{\delta,\delta'}
 \end{array}$$

indeed, for a suitable representative h_Q of $\tilde{h}_{\delta}^{\delta,\delta'}(Q)$, it follows from the diagram above that we have

$$h_Q(1)e_{\mu}^{\delta,\delta'}(1) = h_Q(1) = e_{\mu}^{\delta,\delta'}(1)h_Q(1)$$

which proves the existence of the left-hand embeddings.

On the other hand, it follows from Proposition 12.12 in [6] that we have a unique isomorphism of $\mathfrak{D}R$ -interior algebras

$$d_{R,\tau}^Q : \text{Ind}_{\theta}(\text{Res}_{\tilde{T}}^{\tilde{Q}}(\ddot{C}_{\delta,\delta'})) \cong \text{Res}_R^Q(\text{Ind}_{\tau}(\ddot{C}_{\delta,\delta'}))$$

mapping $1 \otimes a$ on $1 \otimes a$ for any $a \in \ddot{C}_{\delta,\delta'}$ such that $\text{Ker}(\tau)$ fixes $1 \otimes a$ in $k \otimes_{k \text{Ker}(\tau)} \ddot{C}_{\delta,\delta'}$; now, consider an orthogonal primitive idempotent decomposition \tilde{L} of the unity element in $\mathbb{C}_{\circ}(\dot{C}_{\delta,\delta'})^{\tilde{T}}$, so that setting

$$\hat{\ell} = 1 \otimes \tilde{\ell} \quad \text{and} \quad \bar{\ell} = \text{Br}_R^{\text{Ind}(\ddot{C}_{\delta,\delta'})}(\hat{\ell}),$$

the family $\{\bar{\ell}\}_{\tilde{\ell} \in \tilde{L}}$ form an orthogonal idempotent decomposition of the unity element in $(\text{Ind}_{\tau}(\ddot{C}_{\delta,\delta'}))(R)$.

Considering $\hat{j} \in \hat{\varepsilon}$, we can choose a representative h_R of $\tilde{h}_{\delta}^{\delta,\delta'}(R)$ such that $h_R(\text{Br}_R^{\hat{A}_{\delta}}(\hat{j}))$ centralizes that family and, since this idempotent is primitive, there is $\tilde{\ell} \in \tilde{L}$ fulfilling

$$h_R(\text{Br}_R^{\hat{A}_{\delta}}(\hat{j}))\bar{\ell} = h_R(\text{Br}_R^{\hat{A}_{\delta}}(\hat{j})) = \bar{\ell}h_R(\text{Br}_R^{\hat{A}_{\delta}}(\hat{j}));$$

let $\tilde{\eta}$ be the point of \tilde{T} on $\mathbb{C}_{\circ}(\ddot{C}_{\delta,\delta'})$ determined by $\tilde{\ell}$; setting $\ddot{C}_{\tilde{\eta}} = (\ddot{C}_{\delta,\delta'})_{\tilde{\eta}}$, the canonical exoembedding $\tilde{f}_{\tilde{\eta}}^{\delta,\delta'} : C_{\tilde{\eta}} \rightarrow \text{Res}_{\tilde{T}}^{\tilde{Q}}(\ddot{C}_{\delta,\delta'})$ determines the following commutative diagram of \mathfrak{D} -algebra exoembeddings

$$\begin{array}{ccc}
 \hat{A}_{\delta}(R) & \xrightarrow{\tilde{h}_{\delta}^{\delta,\delta'}(R)} & (\text{Ind}_{\tau}(\ddot{C}_{\delta,\delta'}))(R) \\
 \uparrow \tilde{f}_{\varepsilon}^{\delta}(R) & & \uparrow (\text{Ind}_{\tau}(\tilde{f}_{\tilde{\eta}}^{\delta,\delta'}))(R) \\
 \hat{A}_{\hat{\varepsilon}}(R) & \longrightarrow & (\text{Ind}_{\theta}(\ddot{C}_{\tilde{\eta}}))(R)
 \end{array} \tag{2.14.1}$$

Thus, applying again Theorem 13.9 in [6] and denoting by $\{\tilde{e}_{\tilde{\nu}}^{\tilde{\eta}}\}_{\tilde{\nu} \in \tilde{S}_{\theta}}$ the corresponding family of exoembeddings to $(\text{Ind}_{\theta}(\ddot{C}_{\tilde{\eta}}))(R)$, from Proposition 13.17 in [6] we get the following commutative

diagram of \mathfrak{D} -algebra exoembeddings

$$\begin{array}{ccc}
 \hat{A}_{\delta}(R) & \xrightarrow{\tilde{h}_{\delta}^{\ddot{\varepsilon}, \delta'}(R)} & (\text{Ind}_{\tau}(\ddot{C}_{\delta, \delta'}))(R) \\
 \uparrow \tilde{f}_{\varepsilon}^{\delta} & \searrow & \nearrow \tilde{e}_{\nu}^{\ddot{\varepsilon}, \delta'} \\
 & \text{Ind}_{\tau_{\nu}}(\ddot{C}_{\delta, \delta'}(R^{\nu})) & \\
 \hat{A}_{\varepsilon}(R) & \xrightarrow{\quad} & (\text{Ind}_{\theta}(\ddot{C}_{\eta}))(R) \\
 & \nearrow & \uparrow (\text{Ind}_{\theta}(\tilde{f}_{\eta}^{\ddot{\varepsilon}, \delta'}))(R) \\
 & \text{Ind}_{\theta_{\nu}}(\ddot{C}_{\eta}(R^{\nu})) & \nearrow \tilde{e}_{\nu}^{\ddot{\eta}}
 \end{array} \quad (2.14.2)$$

where θ_{ν} denotes the corresponding restriction of θ .

But, recall that any local point $\ddot{\varepsilon} \times \varepsilon'$ of a subgroup \ddot{R} of \ddot{Q} over $\ddot{C}_{\delta, \delta'}$ is given by a local point $\ddot{\varepsilon}$ over $\mathbb{C}_{\circ}(\hat{A}_{\delta})$ and a local point ε' over $(kG')_{\delta'}$ [4, Proposition 5.6], and that, setting $R' = \tau'(\ddot{R})$ and denoting by $\rho: \ddot{R} \rightarrow R$ and $\rho': \ddot{R} \rightarrow R'$ the respective restrictions of θ and θ' , we have a commutative diagram of $\mathfrak{D}\ddot{R}$ -algebra exoembeddings [6, 14.2]

$$\begin{array}{ccc}
 & \text{Res}_{\ddot{R}}^{\ddot{Q}}(\ddot{C}_{\delta, \delta'}) & \\
 \nearrow \tilde{f}_{\ddot{\varepsilon}, \varepsilon'}^{\ddot{\varepsilon}, \delta'} & & \nwarrow \tilde{f}_{\ddot{\varepsilon}}^{\ddot{\varepsilon}} \otimes \tilde{f}_{\varepsilon'}^{\delta'} \\
 (\ddot{C}_{\delta, \delta'})_{\ddot{\varepsilon} \times \varepsilon'} & \xrightarrow{\quad} & C_{\ddot{\varepsilon}, \varepsilon'} = \hat{A}_{\ddot{\varepsilon}} \otimes \text{Res}_{\rho'}(kG')_{\varepsilon'}
 \end{array} \quad (2.14.3)$$

moreover, choosing a defect pointed group $\ddot{R}_{\ddot{\varepsilon}}$ of \ddot{T}_{η} , the *Green Indecomposability Theorem* suitably generalized [6, 2.12.2] guarantees that

$$\ddot{C}_{\eta} \cong \text{Ind}_{\ddot{R}_{\ddot{\varepsilon}}}^{\ddot{T}}((\ddot{C}_{\delta, \delta'})_{\ddot{\varepsilon} \times \varepsilon'}). \quad (2.14.4)$$

Then, this isomorphism and the commutative diagrams (2.14.1) and (2.14.3) provide the bottom part of the following commutative diagram of \mathfrak{D} -algebra exoembeddings

$$\begin{array}{ccc}
 & \hat{A}(R) & \\
 \nearrow \tilde{f}_{\delta}^{\delta} & & \nwarrow \tilde{g}_{\delta, \delta'}^{\delta} \\
 \hat{A}_{\delta}(R) & \xrightarrow{\tilde{h}_{\delta}^{\ddot{\varepsilon}, \delta'}(R)} & (\text{Ind}_{\tau}(\ddot{C}_{\delta, \delta'}))(R) \\
 \uparrow \tilde{f}_{\varepsilon}^{\delta} & & \uparrow \tilde{g}_{\varepsilon, \varepsilon'}^{\ddot{\varepsilon}, \delta'}(R) \\
 \hat{A}_{\varepsilon}(R) & \xrightarrow{\tilde{h}_{\varepsilon}^{\ddot{\varepsilon}, \varepsilon'}(R)} & (\text{Ind}_{\rho}(\ddot{C}_{\varepsilon, \varepsilon'}))(R)
 \end{array}$$

Firstly note that the bottom embedding proves that R has a local point over $\text{Ind}_{\rho}(\ddot{C}_{\varepsilon, \varepsilon'})$ and therefore we necessarily have $\rho(\ddot{R}) = R$; thus, this commutative diagram proves that $(R_{\hat{\varepsilon}}, \ddot{R}_{\hat{\varepsilon}}, R'_{\varepsilon'})$ is a local

tracing triple on \hat{A} , \check{A} and kG' (cf. 2.9), and that it is contained in $(Q_{\delta}, \check{Q}_{\delta}, Q'_{\delta'})$ (cf. 2.10). Moreover, we have [6, Corollary 12.7]

$$\text{Ind}_{\theta}(\check{C}_{\check{\eta}}) \cong \text{Ind}_{\theta}(\text{Ind}_{\check{R}}^{\check{\tau}}((\check{C}_{\check{\delta}, \delta'})^{\check{\varepsilon} \times \varepsilon'})) \cong \text{Ind}_{\rho}((\check{C}_{\check{\delta}, \delta'})^{\check{\varepsilon} \times \varepsilon'});$$

but, applying again Theorem 13.9 in [6] and denoting by $\{\tilde{e}_{\omega}^{\check{\varepsilon} \times \varepsilon'}\}_{\omega \in \check{S}_{\rho}}$ the corresponding family of pairwise of exoembeddings to $(\text{Ind}_{\rho}((\check{C}_{\check{\delta}, \delta'})^{\check{\varepsilon} \times \varepsilon'}))(R)$, we get two families of embeddings to this \mathfrak{D} -algebra, namely $\{\tilde{e}_{\omega}^{\check{\varepsilon} \times \varepsilon'}\}_{\omega \in \check{S}_{\rho}}$ and the composition of $\{\tilde{e}_{\check{\nu}}^{\check{\eta}}\}_{\check{\nu} \in \check{S}_{\theta}}$ with the isomorphism above, both providing orthogonal idempotent decompositions of the unity element; note that $\tilde{e}_{\check{\nu}}^{\check{\eta}}(1) \neq 0$ forces $\check{C}_{\check{\eta}}(R^{\check{\nu}}) \neq \{0\}$, so that a \check{T} -conjugate of $R^{\check{\nu}}$ is contained in \check{R} [6, 2.9.5]. Then, it follows from the uniqueness part of this theorem that, for some section $\omega \in \check{S}_{\rho}$, $\tilde{e}_{\check{\nu}}^{\check{\eta}}$ coincides with $\tilde{e}_{\omega}^{\check{\varepsilon} \times \varepsilon'}$, so that a $\text{Ker}(\theta)$ -conjugate of ν coincides with ω ; actually, up to modifying our choice of the defect pointed group $\check{R}_{\check{\varepsilon}}$, we may assume that ν coincides with ω .

Finally, denoting by ρ_{ν} the corresponding restriction of ρ , from the bottom exoembedding in diagram (2.14.3) and from the isomorphism above we get a \mathfrak{D} -algebra exoembedding

$$\tilde{c} : (\text{Ind}_{\theta}(\check{C}_{\check{\eta}}))(R) \rightarrow (\text{Ind}_{\rho}(\check{C}_{\check{\varepsilon}, \varepsilon'}))(R)$$

and it follows from Proposition 13.17 in [6] applied to the bottom exoembedding in diagram (2.14.3) that, for a representative c of \tilde{c} , we have

$$c(e_{\check{\nu}}^{\check{\eta}}(1))e_{\check{\nu}}^{\check{\varepsilon}, \varepsilon'}(1) = c(e_{\check{\nu}}^{\check{\eta}}(1)) = e_{\check{\nu}}^{\check{\varepsilon}, \varepsilon'}(1)c(e_{\check{\nu}}^{\check{\eta}}(1)).$$

At this point, from diagram (2.14.2) above and from this equality we get the following commutative diagram of \mathfrak{D} -algebra exoembeddings

$$\begin{array}{ccccc} \hat{A}_{\hat{\delta}}(R) & \xrightarrow{\tilde{h}_{\hat{\delta}}^{\check{\delta}, \delta'}(R)} & & (\text{Ind}_{\tau}(\check{C}_{\check{\delta}, \delta'}))(R) & \\ \uparrow \tilde{f}_{\varepsilon}(R) & \searrow & \text{Ind}_{\tau_{\nu}}(\check{C}_{\check{\delta}, \delta'}(R^{\nu})) & \nearrow e_{\check{\nu}}^{\check{\delta}, \delta'} & \\ & & & & \uparrow \tilde{g}_{\check{\varepsilon}, \varepsilon'}^{\check{\delta}, \delta'}(R) \\ \hat{A}_{\hat{\varepsilon}}(R) & \xrightarrow{\tilde{h}_{\hat{\varepsilon}}^{\check{\varepsilon}, \varepsilon'}(R)} & & (\text{Ind}_{\rho}(\check{C}_{\check{\varepsilon}, \varepsilon'}))(R) & \\ & \searrow & \text{Ind}_{\rho_{\nu}}(\check{C}_{\check{\varepsilon}, \varepsilon'}(R^{\nu})) & \nearrow e_{\check{\nu}}^{\check{\varepsilon}, \varepsilon'} & \end{array}$$

which proves that ν is a section of $(R_{\hat{\varepsilon}}, \check{R}_{\check{\varepsilon}}, R'_{\varepsilon'})$; indeed, choosing suitable representatives f of $\tilde{f}_{\varepsilon}(R)$ and g of $\tilde{g}_{\check{\varepsilon}, \varepsilon'}^{\check{\delta}, \delta'}(R)$, we may assume that the idempotents $h_R(f(1))$, $e_{\check{\nu}}^{\check{\delta}, \delta'}(1)$ and $g(1)$ centralize each other; then, we have

$$h_R(f(1))e_{\check{\nu}}^{\check{\delta}, \delta'}(1)g(1) = h_R(f(1))$$

and this equality proves the existence of the bottom left-hand exoembedding. We are done. \square

3. The extended Brauer quotient

3.1. Here we extend to our new setting our construction in [7, Section 3]. Let G be a finite group and V be a kG -module; recall that for any subgroup P of G , we denote by V^P the k -submodule of all P -fixed elements of V , by $V(P)$ the quotient

$$V(P) = V^P / \sum_R V_R^P,$$

where R runs over the set of all proper subgroups of P and V_R^P is the image of the usual relative trace map $\text{Tr}_R^P : V^R \rightarrow V^P$, and by Br_P^V the canonical surjective homomorphism $V^P \rightarrow V(P)$, which is the so-called *Brauer homomorphism* associated to P and V . Obviously, the kG -module structure on V induces $kN_G(P)$ -module structures on both V^P and $V(P)$, and Br_P^V is a homomorphism of $kN_G(P)$ -modules. If H is normal in G and $V = kH$, then for any p -subgroup P of G , P acts on H by conjugation and thus on kH and it is easily checked that $V(P) \cong kC_H(P)$ as $kN_G(P)$ -modules; in this case, we often identify $V(P)$ with $kC_H(P)$. If V' is a p -permutation kG -module, there is a $kN_G(P)$ -module isomorphism [6, Lemma 7.10]

$$\text{Br}_P^{V, V'} : (V \otimes_k V')(P) \cong V(P) \otimes_k V'(P)$$

mapping $\text{Br}_P^{V \otimes_k V'}(v \otimes v')$ onto $\text{Br}_P^V(v) \otimes \text{Br}_P^{V'}(v')$ for any $v \in V^P$ and $v' \in V'^P$. If $f : V \rightarrow V''$ is a homomorphism of kG -modules, for any subgroup P of G and any subgroup R of P , we have $f(V^P) \subset V''^P$ and $f(V_R^P) \subset V_R''^P$, so that f induces a $kN_G(P)$ -module homomorphism $f(P) : V(P) \rightarrow V''(P)$.

3.2. Let \ddot{M} be a $\mathfrak{D}(G \times G)$ -module, Q a p -subgroup of G and K a subgroup of the automorphism group $\text{Aut}(Q)$ of Q . For any $\varphi \in K$, as above we denote by $\Delta_\varphi : Q \rightarrow Q \times Q$ the φ -twisted diagonal homomorphism mapping $u \in Q$ onto $(\varphi(u), u)$ and we set

$$N_{\ddot{M}}^\varphi(Q) = \ddot{M}^{\Delta_\varphi(Q)} \quad \text{and} \quad N_{\ddot{M}}^K(Q) = \bigoplus_{\varphi \in K} N_{\ddot{M}}^\varphi(Q).$$

It is clear that $N_{\ddot{M}}^\varphi(Q)$ inherits a \mathfrak{D} -module structure and, for any elements $x, y \in N_G^K(Q)$, denoting by \bar{x} and \bar{y} their images in $\text{Aut}(Q)$, we have

$$(x, y) \cdot N_{\ddot{M}}^\varphi(Q) = N_{\ddot{M}}^{\bar{x} \circ \varphi \circ \bar{y}}(Q);$$

that is to say, the $\mathfrak{D}(G \times G)$ -module structure over \ddot{M} determines over $N_{\ddot{M}}^K(Q)$ a $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module structure.

3.3. Similarly, for any subgroup R of Q , it is clear that the relative trace map from $\ddot{M}^{\Delta_\varphi(R)}$ to $\ddot{M}^{\Delta_\varphi(Q)}$ is a \mathfrak{D} -module homomorphism and that we get

$$(x, y) \cdot \ddot{M}_{\Delta_\varphi(R)}^{\Delta_\varphi(Q)} = \ddot{M}_{\Delta_{\bar{x} \circ \varphi \circ \bar{y}}(R)}^{\Delta_{\bar{x} \circ \varphi \circ \bar{y}}(Q)};$$

thus, the direct sum $\bigoplus_{\varphi \in K} \text{Ker}(\text{Br}_{\Delta_\varphi(Q)}^{\ddot{M}})$ is a $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -submodule of $N_{\ddot{M}}^K(Q)$. As in [7], in the present paper we are mainly interested in the following quotient

$$\bar{N}_{\ddot{M}}^K(Q) = N_{\ddot{M}}^K(Q) / \left(\bigoplus_{\varphi \in K} \text{Ker}(\text{Br}_{\Delta_\varphi(Q)}^{\ddot{M}}) \right) = \bigoplus_{\varphi \in K} \ddot{M}(\Delta_\varphi(Q))$$

– called the *extended Brauer quotient* of \dot{M} at (Q, K) . If $f: \dot{M} \rightarrow \dot{M}'$ is a $\mathfrak{D}(G \times G)$ -module homomorphism, it is clear that f maps $N_{\dot{M}}^{\varphi}(Q)$ inside $N_{\dot{M}'}^{\varphi}(Q)$ and induces a $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module homomorphism $N_f^K(Q): N_{\dot{M}}^K(Q) \rightarrow N_{\dot{M}'}^K(Q)$ such that

$$(N_f^K(Q)) \left(\bigoplus_{\varphi \in K} \text{Ker}(\text{Br}_{\Delta_{\varphi}(Q)}^{\dot{M}}) \right) \subset \bigoplus_{\varphi \in K} \text{Ker}(\text{Br}_{\Delta_{\varphi}(Q)}^{\dot{M}'});$$

thus, f induces a $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module homomorphism

$$\tilde{N}_f^K(Q): \tilde{N}_{\dot{M}}^K(Q) \rightarrow \tilde{N}_{\dot{M}'}^K(Q).$$

3.4. If A is a $\mathfrak{D}G$ -interior algebra, A naturally admits a $\mathfrak{D}(G \times G)$ -module structure, so that the extended Brauer quotient $\tilde{N}_A^K(Q)$ makes sense, and then $\tilde{N}_A^K(Q)$ becomes a $\mathfrak{D}N_G^K(Q)$ -interior algebra with the distributive product defined by the equality [7, 3.2]

$$\text{Br}_{\Delta_{\varphi}(Q)}^A(a) \text{Br}_{\Delta_{\varphi'}(Q)}^A(a') = \text{Br}_{\Delta_{\varphi \circ \varphi'}(Q)}^A(aa')$$

for any $\varphi, \varphi' \in K$, any $a \in N_A^{\varphi}(Q)$ and any $a' \in N_A^{\varphi'}(Q)$, and with the k -algebra homomorphism

$$\mathfrak{D}N_G^K(Q) \rightarrow \tilde{N}_A^K(Q)$$

mapping $z \in \mathfrak{D}$ onto $\text{Br}_{\Delta_{\text{id}_Q}(Q)}^A(z \cdot 1_A)$ and $x \in N_G^K(Q)$ onto $\text{Br}_{\Delta_{\bar{x}}(Q)}^A(x \cdot 1_A)$.

3.5. Some general results in [7, Section 3], as Proposition 3.4, can be easily extended to the new context, but Propositions 3.8 and 3.9 deserve a careful analysis.² Let P be a p -subgroup of G containing Q and S a $\mathfrak{D}P$ -interior algebra which is a matrix algebra over k and admits a P -stable basis; according to Corollary 5.8 in [4], $S(Q)$ is a matrix algebra over k too (possibly zero!); in particular, $S(Q)$ becomes a $\mathfrak{D}N_P^K(Q)$ -interior algebra and at most Q has one local point over the kP -interior algebra S . On the other hand, the canonical homomorphism $N_P^K(Q) \rightarrow K$ and the trivial \mathfrak{D} -interior structure induces a $\mathfrak{D}N_P^K(Q)$ -interior algebra structure on kK .

Lemma 3.6. *With the notation and the hypothesis above, assume that Q has a local point χ over the kP -interior algebra S and that the kQ -interior algebra S_{χ} is K -stable. Then, we have a $\mathfrak{D}N_P^K(Q)$ -interior algebra isomorphism*

$$\tilde{N}_S^K(Q) \cong S(Q) \otimes_k kK$$

compatible with the K -gradings.

Proof. Since $S(Q)$ is a full matrix algebra over k and a unitary subalgebra of $\tilde{N}_S^K(Q)$, it follows from Proposition 2.1 in [2] that the product induces a k -algebra isomorphism

$$S(Q) \otimes_k C_{\tilde{N}_S^K(Q)}(S(Q)) \cong \tilde{N}_S^K(Q). \quad (3.6.1)$$

² Actually, the fact that Q need not be contained in the K -normalizer of Q has been occasionally forgotten in [7] but the reader will easily modify the concerned statements.

For any $x \in N_p^K(Q)$, the element $\text{Br}_{\Delta_{\bar{x}}(Q)}^S(x \cdot 1_S) \in \bar{N}_S^K(Q)$ obviously induces on $S(Q)$ the same automorphism as $x \cdot 1_{S(Q)} \in S(Q)$; thus, the element $x^{-1} \cdot \text{Br}_{\Delta_{\bar{x}}(Q)}^S(x \cdot 1_S)$ belongs to $C_{\bar{N}_S^K(Q)}(S(Q))$ and it is easily checked that the correspondence mapping $x \in N_p^K(Q)$ onto $x^{-1} \cdot \text{Br}_{\Delta_{\bar{x}}(Q)}^S(x \cdot 1_S)$ defines a group homomorphism

$$N_p^K(Q) \rightarrow C_{\bar{N}_S^K(Q)}(S(Q))^*.$$

Then, endowed with this group homomorphism and with the trivial \mathfrak{D} -interior algebra structure, $C_{\bar{N}_S^K(Q)}(S(Q))$ becomes a $\mathfrak{D}N_p^K(Q)$ -interior algebra and isomorphism (3.6.1) a $\mathfrak{D}N_p^K(Q)$ -interior algebra isomorphism.

Now, in order to complete the proof, it suffices to show that there exists a $kN_p^K(Q)$ -interior algebra isomorphism

$$kK \cong C_{\bar{N}_S^K(Q)}(S(Q)) \quad (3.6.2)$$

compatible with the K -gradings; actually, since $N_p^K(Q)$ is a p -group and all the terms in the gradings have dimension one, it suffices to exhibit a simple k -algebra isomorphism. Choose $\ell \in \chi$ and set $S_\chi = \ell S\ell$; it follows from Proposition 3.4 in [7] that we have a k -algebra embedding

$$\bar{f} : \bar{N}_{S_\chi}^K(Q) \rightarrow \bar{N}_S^K(Q);$$

since $\bar{f}(\ell)$ is a primitive idempotent of $S(Q)$, it follows from isomorphism (3.6.1) that \bar{f} induces a k -algebra isomorphism

$$\bar{N}_{S_\chi}^K(Q) \cong C_{\bar{N}_S^K(Q)}(S(Q))$$

compatible with the K -gradings.

But, to assume that S_χ is K -stable amounts to say that, for any $\varphi \in K$, there is $a_\varphi \in (S_\chi)^*$ such that $a_\varphi \cdot u \cdot a_\varphi^{-1} = \varphi(u)$ for any $u \in Q$ and therefore a_φ belongs to $N_{S_\chi}^\varphi(Q)$; thus, since $S_\chi(Q) \cong k$, we get $\bar{N}_{S_\chi}^\varphi(Q) = k \cdot \bar{a}_\varphi$ and isomorphism (3.6.2) follows from Dade's Theorem in [3]. \square

Proposition 3.7. *With the notation and the hypothesis above, assume that Q has a local point χ over the kP -interior algebra S and that the kQ -interior algebra S_χ is K -stable. Then, for any $\mathfrak{D}P$ -interior algebra B we have a $\mathfrak{D}N_p^K(Q)$ -interior algebra isomorphism*

$$\bar{N}_{S \otimes_k B}^K(Q) \cong S(Q) \otimes_k \bar{N}_B^K(Q)$$

compatible with the K -gradings.

Proof. For any $\varphi \in K$, since there is $a_\varphi \in (S_\chi)^*$ such that $a_\varphi \cdot u \cdot a_\varphi^{-1} = \varphi(u)$ for any $u \in Q$, S_χ still admits a $\Delta_\varphi(Q)$ -stable basis; moreover, according to [5, 3.13], there is a permutation kQ -module V such that we have a kQ -interior algebra embedding $\text{Res}_Q^P(S) \rightarrow \text{End}_k(V) \otimes_k S_\chi$ and therefore S admits a $\Delta_\varphi(Q)$ -stable basis too. Hence, we have an isomorphism

$$\bar{N}_{S \otimes_k B}^K(Q) \cong \bigoplus_{\varphi \in K} S(\Delta_\varphi(Q)) \otimes_k B(\Delta_\varphi(Q)).$$

But, it is clear that the direct sum

$$\bigoplus_{\varphi \in K} S(\Delta_{\varphi}(Q)) \otimes_k B(\Delta_{\varphi}(Q)) = \bigoplus_{\varphi \in K} \bar{N}_S^{\varphi}(Q) \otimes_k \bar{N}_B^{\varphi}(Q)$$

is a $\mathcal{D}N_p^K(Q)$ -interior subalgebra of $\bar{N}_S^K(Q) \otimes_k \bar{N}_B^K(Q)$; moreover, it follows from Lemma 3.6 above that we have

$$\bar{N}_S^{\varphi}(Q) = S(Q)\bar{s}_{\varphi} = \bar{s}_{\varphi}S(Q)$$

for suitable elements $s_{\varphi} \in N_S^{\varphi}(Q)$ fulfilling $\bar{s}_{\varphi}\bar{s}_{\varphi'} = \bar{s}_{\varphi \circ \varphi'}$ and $s_{\bar{x}} = x \cdot 1_S$ for any $x \in N_p^K(Q)$. Now, it is easy to build the announced isomorphism. \square

3.8. As in [7], the K -extended Brauer quotient $\bar{N}_A^K(Q)$ will replace the ordinary Brauer quotient $A(Q)$ and, coherently, we will extend to $\bar{N}_A^K(Q)$ Theorem 13.9 in [6]; thus, let $\rho: G \rightarrow \bar{G}$ be a surjective group homomorphism of kernel W and \bar{Q} a p -subgroup of \bar{G} , and assume that A is projective as $k(W \times W)$ -module and that a Sylow p -subgroup of $G \times_{\bar{G}} G$ stabilizes a basis of A . It is clear that $A^W(\bar{Q})$ has a $\mathcal{D}C_G(\rho^{-1}(\bar{Q}))$ -interior algebra structure; similarly, denoting by \mathcal{C} the (possibly empty!) set of all the complements of W in $\rho^{-1}(\bar{Q})$, W acts on \mathcal{C} and $(\prod_{Q \in \mathcal{C}} A(Q))^W$ also becomes a $\mathcal{D}C_G(\rho^{-1}(\bar{Q}))$ -interior algebra. Then, Lemma 13.6 in [6] states that the inclusion $A^W \subset A$ induces a $\mathcal{D}C_G(\rho^{-1}(\bar{Q}))$ -interior algebra isomorphism

$$A^W(\bar{Q}) \cong \left(\prod_{Q \in \mathcal{C}} A(Q) \right)^W. \quad (3.8.1)$$

Thus, if $A^W(\bar{Q}) \neq \{0\}$ then $\mathcal{C} \neq \emptyset$; in this case, for any complement Q of W in $\rho^{-1}(\bar{Q})$, we denote by

$$c_{\rho, Q}(A) : A(Q)^{C_W(Q)} \rightarrow A^W(\bar{Q}) \quad (3.8.2)$$

the $\mathcal{D}C_G(\rho^{-1}(\bar{Q}))$ -interior algebra embedding induced by isomorphism (3.8.1).

Proposition 3.9. *With the notation and the hypothesis above, let \bar{K} be a subgroup of $\text{Aut}(\bar{Q})$ and Q a complement of W in $\rho^{-1}(\bar{Q})$, and denote by K the image of \bar{K} in $\text{Aut}(Q)$ and by $\rho_Q^K : N_Q^K(Q) \rightarrow N_{\bar{G}}^{\bar{K}}(\bar{Q})$ the group homomorphism both determined by ρ . Then there exists a $\mathcal{D}\rho(N_Q^K(Q))$ -interior algebra embedding*

$$e_{\rho, Q}^K(A) : \text{Ind}_{\rho_Q^K}(\bar{N}_A^K(Q)) \rightarrow \bar{N}_{\text{Ind}_{\rho}(A)}^{\bar{K}}(\bar{Q})$$

which is compatible with the K - and \bar{K} -gradings and makes commutative the following diagram

$$\begin{array}{ccc} \text{Ind}_{\rho_Q^K}(\bar{N}_A^K(Q)) & \xrightarrow{e_{\rho, Q}^K(A)} & \bar{N}_{\text{Ind}_{\rho}(A)}^{\bar{K}}(\bar{Q}) \\ d_{\rho_Q^K}(A(Q)) \uparrow & & \uparrow (d_{\rho}(A))(\bar{Q}) \\ A(Q)^{C_W(Q)} & \xrightarrow{c_{\rho, Q}(A)} & A^W(\bar{Q}) \end{array} \quad (3.9.1)$$

Proof.³ Since a Sylow p -subgroup of $G \times_{\bar{G}} G$ stabilizes a basis of A and, considered as $k(W \times W)$ -module, A is projective, it is easy to check that $A(Q)$ is also projective as $k(C_W(Q) \times C_W(Q))$ -module and therefore $A^{W \cdot Q}$ maps onto $A(Q)^{C_W(Q)}$; in particular, the following commutative diagram determines $c_{\rho, Q}(A)$

$$\begin{array}{ccc} A(Q)^{C_W(Q)} & \xrightarrow{c_{\rho, Q}(A)} & A^W(\bar{Q}) \\ & \nwarrow \quad \nearrow & \\ & A^{W \cdot Q} & \end{array}$$

More generally, denoting by \bar{N} the image of $Q \cdot N_G^K(Q)$ in $\text{Aut}(W)$, it is clear that $N_A^{\bar{N}}(W)$ can be identified to a $\mathfrak{D}(Q \cdot N_G^K(Q))$ -interior subalgebra of A (cf. 2.6) and that $N_{N_A^{\bar{N}}(W)}^K(Q)$ contains $A^{W \cdot Q}$; moreover, denoting by $\bar{\bar{N}}$ the image of \bar{N} in $\text{Aut}(C_W(Q))$ we claim that

$$N_{N_A^{\bar{N}}(W)}^K(Q) \subset N_{N_A^{\bar{N}}(Q)}^{\bar{\bar{N}}}(C_W(Q));$$

indeed, for any $\varphi \in K$, it is quite clear that any element in $N_A^{\bar{N}}(W)^{\Delta_{\varphi}(Q)}$ still belongs to $N_A^{\bar{N}}(C_W(Q))^{\Delta_{\varphi}(Q)}$ and that

$$N_A^{\bar{N}}(C_W(Q))^{\Delta_{\varphi}(Q)} = \bigoplus_{\sigma \in \bar{\bar{N}}} A^{\Delta_{\sigma}(C_W(Q)) \cdot \Delta_{\varphi}(Q)} \subset N_{N_A^{\bar{N}}(Q)}^{\bar{\bar{N}}}(C_W(Q)).$$

Hence, we have the unitary $\mathfrak{D}(N_G^K(Q))$ -interior algebra homomorphism determined by $d_{\rho_Q^K}(\bar{N}_A^K(Q))$ (cf. 2.6)

$$N_{N_A^{\bar{N}}(W)}^K(Q) \subset N_{N_A^{\bar{N}}(Q)}^{\bar{\bar{N}}}(C_W(Q)) \rightarrow N_{\bar{N}_A^K(Q)}^{\bar{\bar{N}}}(C_W(Q)) \rightarrow \text{Ind}_{\rho_Q^K}(\bar{N}_A^K(Q)).$$

Similarly, $d_{\rho}(A)$ induces a unitary $\mathfrak{D}(Q \cdot N_G^K(Q))$ -interior algebra homomorphism

$$N_A^{\bar{N}}(W) \rightarrow \text{Ind}_{\rho}(A)$$

and therefore we still get a unitary $\mathfrak{D}(N_G^K(Q))$ -interior algebra homomorphism

$$N_{N_A^{\bar{N}}(W)}^K(Q) \rightarrow \bar{N}_{N_A^{\bar{N}}(W)}^K(Q) \rightarrow \bar{N}_{\text{Ind}_{\rho}(A)}^{\bar{K}}(\bar{Q}).$$

On the other hand, since a Sylow p -subgroup of $W \cdot Q$ stabilizes a basis of A which is projective as kW -module, there is an idempotent i_Q of $A_Q^{W \cdot Q}$ lifting the image $(c_{\rho, Q}(A))(1)$ of the unity element of $A(Q)$ (cf. (3.8.2)); then, setting $B = i_Q A i_Q$, B remains a $k(W \cdot Q)$ -interior algebra, and the embedding determined by the inclusion $B \subset A$ induces a $k(C_W(Q) \cdot Q)$ -interior algebra isomorphism $\bar{N}_B^K(Q) \cong \bar{N}_A^K(Q)$ and two kQ -interior algebra embeddings

³ It is an extension of the proof of Theorem 13.9 in [6] from the Brauer quotient to the extended Brauer quotient, except that in [6] the proof of the compatibility with the centralizer interior structures is missed.

$$N_B^{\bar{N}}(W) \rightarrow N_A^{\bar{N}}(W) \quad \text{and} \quad \text{Ind}_\rho(B) \rightarrow \text{Ind}_\rho(A)$$

where we still denote by ρ its restriction to $W \cdot Q$.

At this point, since the following diagram

$$\begin{array}{ccccc} A(Q)^{C_W(Q)} \cong B(Q)^{C_W(Q)} & \cong & B^W(\bar{Q}) & \rightarrow & A^W(\bar{Q}) \\ & \nwarrow & \nearrow & & \\ & B^{W \cdot Q} & & & \end{array}$$

is commutative and also determines $c_{\rho, Q}(A)$, in order to prove the existence of $e_{\rho, Q}^K(A)$, it suffices to prove the existence of a k -algebra isomorphism which makes commutative the following diagram

$$\begin{array}{ccccc} \text{Ind}_{\rho_Q^K}(\bar{N}_A^K(Q)) \cong \text{Ind}_{\rho_Q^K}(\bar{N}_B^K(Q)) & \cong & \bar{N}_{\text{Ind}_\rho(B)}^{\bar{K}}(\bar{Q}) & \rightarrow & \bar{N}_{\text{Ind}_\rho(A)}^{\bar{K}}(\bar{Q}) \\ & \nwarrow & \nearrow & & \\ & N_{N_B^{\bar{N}}(W)}^K(Q) & & & \end{array}$$

where we still denote by ρ_Q^K its restriction to $C_W(Q) \cdot Z(Q)$.

Indeed, the \mathfrak{D} -interior structure of both members of the guessed isomorphism is completely determined by the \mathfrak{D} -interior structure of $N_{N_B^{\bar{N}}(W)}^K(Q)$ since the arrows are \mathfrak{D} -linear; moreover, although $N_{N_B^{\bar{N}}(W)}^K(Q)$ need not be a $kN_G^K(Q)$ -interior subalgebra of $N_{N_A^{\bar{N}}(W)}^K(Q)$, for any $x \in N_G^K(Q)$, the arrows send the element $i_Q \cdot x \cdot i_Q$ of $N_{N_B^{\bar{N}}(W)}^K(Q)$ to

$$1 \otimes \text{Br}_Q(i_Q \cdot x \cdot i_Q) = x \cdot (1 \otimes \text{Br}_Q(1)) \quad \text{and} \quad \text{Br}_{\bar{Q}}(1 \otimes i_Q \cdot x \cdot i_Q) = x \cdot \text{Br}_{\bar{Q}}(1 \otimes i_Q)$$

in $\text{Ind}_{\rho_Q^K}(\bar{N}_A^K(Q))$ and $\bar{N}_{\text{Ind}_\rho(A)}^{\bar{K}}(\bar{Q})$ respectively, since $\text{Br}_Q(i_Q) = 1$ and

$$\text{Br}_{\bar{Q}}(1 \otimes i_Q) = ((d_\rho(A))(\bar{Q}) \circ c_{\rho, Q}(A))(1)$$

which is fixed by the action of $N_G^K(Q)$.

Since $i_Q = 1_B$ belongs to $B_Q^{W \cdot Q}$, it follows from Theorem 2.4 that we have a Higman *exoembedding* of $k(W \cdot Q)$ -interior algebras

$$\tilde{h} : B \rightarrow \text{Ind}_Q^{W \cdot Q}(\text{Res}_Q^{W \cdot Q}(B)) = C$$

which clearly induces respective $k(C_W(Q))$ - and kQ -interior algebra embeddings

$$\bar{N}_B^K(Q) \rightarrow \bar{N}_C^K(Q), \quad \text{Ind}_\rho(B) \rightarrow \text{Ind}_\rho(C) \quad \text{and} \quad N_B^{\bar{N}}(W) \rightarrow N_C^{\bar{N}}(W)$$

and then we get the following commutative diagram

$$\begin{array}{ccc}
 \mathrm{Ind}_{\rho_Q^K}(\bar{N}_B^K(Q)) & \longrightarrow & \mathrm{Ind}_{\rho_Q^K}(\bar{N}_C^K(Q)) \\
 \uparrow & & \uparrow \\
 N_{N_B^{\bar{N}}(W)}^K(Q) & \longrightarrow & N_{N_C^{\bar{N}}(W)}^K(Q) \\
 \downarrow & & \downarrow \\
 \bar{N}_{\mathrm{Ind}_\rho(B)}^{\bar{K}}(\bar{Q}) & \longrightarrow & \bar{N}_{\mathrm{Ind}_\rho(C)}^{\bar{K}}(\bar{Q})
 \end{array}$$

Consequently, since the vertical arrows are unitary, it suffices to show the existence of a k -algebra isomorphism which makes commutative the following diagram

$$\begin{array}{ccc}
 \mathrm{Ind}_{\rho_Q^K}(\bar{N}_C^K(Q)) & \cong & \bar{N}_{\mathrm{Ind}_\rho(C)}^{\bar{K}}(\bar{Q}) \\
 & \nwarrow \quad \nearrow & \\
 N_{N_C^{\bar{N}}(W)}^K(Q) & &
 \end{array} \tag{3.9.2}$$

But, on the one hand, it follows from Corollary 12.7 in [6] that

$$\mathrm{Ind}_\rho(C) = \mathrm{Ind}_\rho(\mathrm{Ind}_Q^{W \cdot Q}(\mathrm{Res}_Q^{W \cdot Q}(B))) \cong \mathrm{Res}_Q^{W \cdot Q}(B). \tag{3.9.3}$$

On the other hand, since we clearly have the orthogonal decomposition

$$1_{\bar{N}_C^K(Q)} = \sum_{x \in C_W(Q)} \mathrm{Br}_Q^C(1 \otimes 1 \otimes 1)^x \tag{3.9.4}$$

by [6, 2.6.4], the embedding $\bar{N}_B^K(Q) \rightarrow \bar{N}_C^K(Q)$ induces an isomorphism of $k(C_W(Q) \cdot Z(Q))$ -interior algebras

$$\mathrm{Ind}_{Z(Q)}^{C_W(Q)Z(Q)}(\bar{N}_B^K(Q)) \cong \bar{N}_C^K(Q)$$

and, applying again Corollary 12.7 in [6], we get

$$\mathrm{Ind}_{\rho_Q^K}(\bar{N}_C^K(Q)) \cong \mathrm{Ind}_{\rho_Q^K}(\mathrm{Ind}_{Z(Q)}^{C_W(Q)Z(Q)}(\bar{N}_B^K(Q))) \cong \bar{N}_B^K(Q). \tag{3.9.5}$$

Finally, it suffices to prove that the images in $\bar{N}_B^K(Q)$ of an element of $N_{N_C^{\bar{N}}(W)}^K(Q)$ through the arrows of diagram (3.9.2) and the isomorphisms (3.9.3) and (3.9.5) coincide. But, it is quite clear that an element a of $N_{N_C^{\bar{N}}(W)}^K(Q)$ has the form

$$a = \sum_{\sigma \in \bar{N}} \sum_{w \in W} \mathrm{Tr}_1^{\Delta_\sigma(W)}(w \otimes a_{\sigma, w} \otimes 1)$$

and its image in $\text{Ind}_\rho(C)$ is equal to

$$1 \otimes a = 1 \otimes \text{Tr}_1^W \left(1 \otimes \sum_{\sigma \in \bar{N}} \sum_{w \in W} a_{\sigma, w} \otimes 1 \right);$$

similarly, if $\varphi \in K$ and a is fixed by $\Delta_\varphi(Q)$, since $\Delta_\varphi(Q)$ stabilizes the following decomposition

$$N_C^{\bar{N}}(W) = \bigoplus_{\sigma \in \bar{N}} \bigoplus_{w \in W} \text{Tr}_1^{\Delta_\sigma(W)}(w \otimes B \otimes 1),$$

the image of $a \in N_C^{\bar{N}}(W)^{\Delta_\varphi(Q)}$ in $\bar{N}_{N_C^{\bar{N}}(W)}^K(Q)$ is equal to

$$\text{Br}_{\Delta_\varphi(Q)}^{N_C^{\bar{N}}(W)}(a) = \text{Br}_{\Delta_\varphi(Q)}^{N_C^{\bar{N}}(W)} \left(\sum_{\sigma \in \bar{N}} \sum_{w \in C_W(Q)} \text{Tr}_1^{\Delta_\sigma(W)}(w \otimes a_{\sigma, w} \otimes 1) \right)$$

and therefore its image in $\bar{N}_{\text{Ind}_\rho(C)}^{\bar{K}}(\bar{Q})$ coincides with

$$\text{Br}_{\Delta_\varphi(Q)}^{\text{Ind}_\rho(C)}(1 \otimes a) = \text{Br}_{\Delta_\varphi(Q)}^{\text{Ind}_\rho(C)} \left(1 \otimes \text{Tr}_1^W \left(1 \otimes \sum_{\sigma \in \bar{N}} \sum_{w \in C_W(Q)} a_{\sigma, w} \otimes 1 \right) \right)$$

which, according to Corollary 12.7 in [6], the isomorphism (3.9.3) maps onto the element $\sum_{\sigma \in \bar{N}} \sum_{w \in C_W(Q)} a_{\sigma, w}$ of $\bar{N}_B^K(Q)$.

On the other hand, since any $\sigma \in \bar{N}$ stabilizes $C_W(Q)$, it follows from decomposition (3.9.4) that the image of $a \in N_C^{\bar{N}}(W)^{\Delta_\varphi(Q)}$ in $\bar{N}_C^K(Q)$ is equal to

$$\text{Br}_{\Delta_\varphi(Q)}^C(a) = \text{Br}_{\Delta_\varphi(Q)}^C \left(\sum_{\sigma \in \bar{N}} \sum_{w, w' \in C_W(Q)} \sigma(w')w \otimes a_{\sigma, w} \otimes w'^{-1} \right)$$

and therefore its image in $\text{Ind}_{\rho_Q}^K(\bar{N}_C^K(Q))$ is equal to

$$1 \otimes \text{Br}_{\Delta_\varphi(Q)}^C(a) = 1 \otimes \text{Tr}_1^{C_W(Q)} \left(1 \otimes \sum_{\sigma \in \bar{N}} \sum_{w \in C_W(Q)} a_{\sigma, w} \otimes 1 \right)$$

which, according to Corollary 12.7 in [6], the isomorphism (3.9.5) maps onto the element $\sum_{\sigma \in \bar{N}} \sum_{w \in C_W(Q)} a_{\sigma, w}$ of $\bar{N}_B^K(Q)$. We are done. \square

Remark 3.10. In the proof above, it is possible to choose the idempotent i_Q , which lifts $(c_{\rho, Q}(A))(1)$ to $A_Q^{W \cdot Q}$, fixed by $N_G^K(Q)$ since the natural homomorphism $A^{W \cdot Q} \rightarrow (A^W)(Q)$ is actually a so-called *covering homomorphism of $N_G(Q)$ -algebras* [4, Section 4] and then it suffices to apply Proposition 4.18 in [4].

In order to prove our claim, we choose a Sylow p -subgroup S of G containing a Sylow p -subgroup of $N_G(Q)$, and then, setting $T = W \cap S$, we choose a $(T \times T) \cdot \Delta(S)$ -stable basis X of A . It is easily checked that the subset $W \otimes X \otimes W$ of $B = \text{Ind}_S^{W \cdot S}(A)$ is a $(W \times W) \cdot \Delta(S)$ -stable basis of B ; from this basis, it is not difficult to compute bases of $B^{W \cdot R}$ and of $(B^W)(Q)^R$ for any subgroup of $N_S(Q)$ containing Q , showing that $B^{W \cdot R}$ maps onto $(B^W)(Q)^R$. Finally, since p does not divide $|(W \cdot S) : S|$, the

canonical $k(W \cdot S \times W \cdot S)$ -module injection $A \rightarrow B$ is split, so that $A^{W \cdot R}$ still maps onto $(A^W)(Q)^R$. We are done.

Lemma 3.11. *If a $\mathfrak{D}G$ -interior algebra A is a 0-split $\mathfrak{D}(G \times G)$ -module then, for any p -subgroup Q of G and any subgroup K of $\text{Aut}(Q)$, the $\mathfrak{D}N_G^K(Q)$ -interior algebra $\bar{N}_A^K(Q)$ is a 0-split $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module and the inclusion $N_{\mathbb{C}_0(A)}^K(Q) \subset \mathbb{C}_0(N_A^K(Q))$ induces a $kN_G^K(P)$ -interior algebra isomorphism*

$$\bar{N}_{\mathbb{H}_0(A)}^K(Q) \cong \mathbb{H}_0(\bar{N}_A^K(Q)).$$

Proof. By the very definition of a 0-split $\mathfrak{D}(G \times G)$ -module (cf. 1.7), there are a $k(G \times G)$ -module M and a contractible $\mathfrak{D}(G \times G)$ -module C such that we have a $\mathfrak{D}(G \times G)$ -module isomorphism

$$A \cong M \oplus C \quad (3.11.1)$$

where we consider the trivial \mathfrak{D} -interior structure on M ; in particular, we have $k(G \times G)$ -module isomorphism

$$\mathbb{C}_0(A) \cong M \oplus \mathbb{C}_0(C) \quad \text{and} \quad \mathbb{B}_0(A) \cong \mathbb{B}_0(C) = \mathbb{C}_0(C)$$

and therefore $\mathbb{H}_0(A) \cong M$ as $\mathfrak{D}(G \times G)$ -modules.

For any p -subgroup Q and any subgroup K of $\text{Aut}(Q)$, isomorphism (3.11.1) induces a $\mathfrak{D}(N_G^K(P) \times N_G^K(P))$ -module isomorphism

$$\bar{N}_A^K(Q) \cong \bar{N}_M^K(Q) \oplus \bar{N}_C^K(Q) \quad (3.11.2)$$

and we claim that $\bar{N}_C^K(Q)$ is a contractible $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module. Indeed, by the very definition of contractibility, there exists $h \in \text{End}_{k(G \times G)}(C)$ such that $\text{id}_C = d \cdot h + h \cdot d$ where $d \in \mathfrak{D}$; but, for any $\varphi \in K$, it is clear that $h(C^{\Delta_\varphi(Q)}) \subset C^{\Delta_\varphi(Q)}$ and therefore h induces a k -endomorphism h_φ of $C^{\Delta_\varphi(Q)}$ which also fulfills $\text{id}_{C^{\Delta_\varphi(Q)}} = d \cdot h_\varphi + h_\varphi \cdot d$; moreover, it is easily checked that $\bigoplus_{\varphi \in K} h_\varphi$ is a $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module endomorphism of $N_C^K(Q)$; hence, $N_C^K(Q)$ is a contractible $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module. Pushing it further, since h is compatible with the relative trace maps, $\bar{N}_C^K(Q)$ is a contractible $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module too.

Consequently, it follows from isomorphism (3.11.2) that $\bar{N}_A^K(Q)$ is a 0-split $\mathfrak{D}(N_G^K(Q) \times N_G^K(Q))$ -module and, applying the argument above, we have a $k(N_G^K(Q) \times N_G^K(Q))$ -module isomorphism $\bar{N}_{\mathbb{H}_0(A)}^K(Q) \cong \mathbb{H}_0(\bar{N}_A^K(Q))$; moreover, it is easily checked that this isomorphism makes commutative the following diagram

$$\begin{array}{ccc} \bar{N}_{\mathbb{H}_0(A)}^K(Q) & \cong & \mathbb{H}_0(\bar{N}_A^K(Q)) \\ \uparrow & & \uparrow \\ N_{\mathbb{C}_0(A)}^K(Q) & \longrightarrow & \mathbb{C}_0(N_A^K(Q)) \end{array}$$

where the vertical arrows are surjective. \square

4. Proof of Theorem 1.12

4.1. Let us come back to our standard setting (cf. 1.8) and assume that kGb and $kG'b'$ are basic Rickard equivalent. Let (P, e) be a maximal Brauer (b, G) -pair and (Q, f) a Brauer (b, G) -pair contained in (P, e) ; note that, for any subgroup H of $N_G(Q, f)$ containing $C_G(Q)$, f is also a block of H .

4.2. As in Theorem 1.12, let K be a subgroup of the image of $N_G(Q, f)$ in $\text{Aut}(Q)$, so that f is also a block of both $N_G^K(Q)$ and $Q \cdot N_G^K(Q)$. Let (R, g) and (T, h) be maximal Brauer $(f, Q \cdot N_G^K(Q))$ - and $(f, Q \cdot C_G(Q))$ -pairs respectively; since R and T contain Q , (R, g) and (T, h) are also Brauer (b, G) -pairs and, without loss of generality, we may assume that

$$(Q, f) \subset (T, h) \subset (R, g) \subset (P, e),$$

so that we clearly have $R = Q \cdot N_P^K(Q)$ and $T = Q \cdot C_P(Q)$; actually, it is easily checked that both (R, g) and (T, h) are *selfcentralizing* (cf. 1.10). Let us denote by γ , ε and ν the local points of P , R and T over kGb associated with e , g and h respectively, so that we have $T_\nu \subset R_\varepsilon \subset P_\gamma$; borrowing our notation from 2.7 above, since we assume that kGb and $kG'b'$ are *Rickard equivalent*, it follows from Proposition 2.8 that γ , ε and ν respectively determine *noncontractile* local points $\hat{\gamma}$, $\hat{\varepsilon}$ and $\hat{\nu}$ of P , R and T over \hat{A} still fulfilling $T_{\hat{\nu}} \subset R_{\hat{\varepsilon}} \subset P_{\hat{\gamma}}$.

4.3. Then, it follows from Theorem 16.15 in [6] that we have a local tracing triple $(P_{\hat{\gamma}}, \ddot{P}_{\hat{\gamma}}, P'_{\gamma'})$; moreover, since we are assuming that the above equivalence is *basic*, each one of the groups $\ddot{P} \times_P \ddot{P}$ and $\ddot{P} \times_{P'} \ddot{P}$ stabilizes a basis of $\text{End}_k(\ddot{N})$ and thus, according to our choice, each of them stabilizes a basis of \hat{S} too (cf. 2.7); then, by [6, 19.3], we know that $(P_{\hat{\gamma}}, \ddot{P}_{\hat{\gamma}}, P'_{\gamma'})$ is *basic* (cf. 2.11). In particular, the group homomorphism $\sigma: \ddot{P} \rightarrow P$ determined by the first projection $G \times G' \rightarrow G$ admits a section and we choose a section $\lambda: P \rightarrow \ddot{P}$ of $(P_{\hat{\gamma}}, \ddot{P}_{\hat{\gamma}}, P'_{\gamma'})$ in the sense of 2.11 above. Moreover, denoting by $\sigma': \ddot{P} \rightarrow P'$ the group homomorphism determined by the second projection $G \times G' \rightarrow G'$ and setting $P^\lambda = \lambda(\ddot{P})$, $\lambda' = \sigma' \circ \lambda$ and $P^{\lambda'} = \lambda'(P)$, the group homomorphism $\lambda': P \rightarrow P'$ is injective since $\ddot{A}_{\hat{\gamma}}(P^\lambda) \neq 0$ [6, 17.4.7]. Thus, by symmetry, λ' is an isomorphism and we have $P^{\lambda'} = P'$,

4.4. At this point, it follows from Theorem 2.14 above, applied to the local pointed group $R_{\hat{\varepsilon}}$ on \hat{A} contained in $P_{\hat{\gamma}}$, that we have a basic local tracing triple $(R_{\hat{\varepsilon}}, \ddot{R}_{\hat{\varepsilon}}, R'_{\varepsilon'})$ contained in $(P_{\hat{\gamma}}, \ddot{P}_{\hat{\gamma}}, P'_{\gamma'})$ and admitting the restriction of λ to R as a section; the same argument applied to the local pointed group $T_{\hat{\nu}}$ contained in $R_{\hat{\varepsilon}}$ and to the local tracing triple $(R_{\hat{\varepsilon}}, \ddot{R}_{\hat{\varepsilon}}, R'_{\varepsilon'})$ provides a basic local tracing triple $(T_{\hat{\nu}}, \ddot{T}_{\hat{\nu}}, T'_{\nu'})$ on \hat{A} , \hat{A} and kG' , such that finally we get the inclusions

$$(T_{\hat{\nu}}, \ddot{T}_{\hat{\nu}}, T'_{\nu'}) \subset (R_{\hat{\varepsilon}}, \ddot{R}_{\hat{\varepsilon}}, R'_{\varepsilon'}) \subset (P_{\hat{\gamma}}, \ddot{P}_{\hat{\gamma}}, P'_{\gamma'})$$

and we know that the restrictions of λ to R and T are respective sections of the triples $(R_{\hat{\varepsilon}}, \ddot{R}_{\hat{\varepsilon}}, R'_{\varepsilon'})$ and $(T_{\hat{\nu}}, \ddot{T}_{\hat{\nu}}, T'_{\nu'})$. Moreover, according to Theorem 18.8 in [6], $P'_{\gamma'}$ is a defect pointed group of b' and, denoting by e' and g' the respective blocks of $C_{G'}(P')$ and $C_{G'}(R')$ determined by γ' and ε' , we have

$$(R', g') \subset (P', e').$$

4.5. We set $R^{\lambda'} = \lambda'(R) \subset R'$, $T^{\lambda'} = \lambda'(T) \subset T'$ and $Q^{\lambda'} = \lambda'(Q) \subset Q'$ (which need not agree with the notation of Theorem 1.12!), and respectively denote by $g^{\lambda'}$, $h^{\lambda'}$ and $f^{\lambda'}$ the blocks of $C_{G'}(R^{\lambda'})$, $C_{G'}(T^{\lambda'})$ and $C_{G'}(Q^{\lambda'})$ fulfilling [6, 2.13.2]

$$(Q^{\lambda'}, f^{\lambda'}) \subset (T^{\lambda'}, h^{\lambda'}) \subset (R^{\lambda'}, g^{\lambda'}) \subset (R', g') \subset (P', e');$$

in particular, since $R_{\hat{\varepsilon}}$ is *selfcentralizing* (cf. 4.2), it follows from Corollary 19.9 in [6] that $R^{\lambda'}$ has a unique local point $\varepsilon^{\lambda'}$ over kG' such that $R^{\lambda'}_{\varepsilon^{\lambda'}} \subset R'_{\varepsilon'}$ and that $R^{\lambda'}_{\varepsilon^{\lambda'}}$ and $R'_{\varepsilon'}$ are both *selfcentralizing* too; then, it follows from Lemma 3.9 in [4], from Theorem 19.7 in [6] and from Lemma 4.18 below that $R^{\lambda'}$ is a defect group of $f^{\lambda'}$ as a block of $Q^{\lambda'} \cdot N_{G'}^{K^{\lambda'}}(Q^{\lambda'})$ and therefore that $N_{P'}^{K^{\lambda'}}(Q^{\lambda'})$ is a defect group of $f^{\lambda'}$ as a block of $N_{G'}^{K^{\lambda'}}(Q^{\lambda'})$.

4.6. Denote by $\rho: \ddot{R} \rightarrow R$ and $\rho': \ddot{R} \rightarrow R'$ the respective group homomorphisms determined by the first and the second projections $G \times G' \rightarrow G$ and $G \times G' \rightarrow G'$; by the very definition of *local tracing triples* (cf. 2.9), there exists a unique $\mathfrak{D}R$ -interior algebra exoembedding

$$\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}: \hat{A}_{\ddot{E}} \rightarrow \text{Ind}_{\rho}(\ddot{A}_{\ddot{E}} \otimes_k \text{Res}_{\rho'}((kG')_{\varepsilon'}))$$

such that we have the corresponding commutative diagram in 2.9; then, setting

$$B_{\ddot{E}, \varepsilon'} = \ddot{A}_{\ddot{E}} \otimes_k \text{Res}_{\rho'}((kG')_{\varepsilon'}),$$

by Proposition 3.4 in [7] suitably extended to \mathfrak{D} -interior algebras, $\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}$ induces a $\mathfrak{D}N_p^K(Q)$ -interior algebra exoembedding which extends $\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}(Q)$

$$\bar{N}_{\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}}^K(Q): \bar{N}_{\hat{A}_{\ddot{E}}}^K(Q) \rightarrow \bar{N}_{\text{Ind}_{\rho}(B_{\ddot{E}, \varepsilon'})}^K(Q).$$

4.7. On the other hand, denoting by K^{λ} the image of K in $\text{Aut}(Q^{\lambda})$ determined by λ , Proposition 3.9 applied to $\pi: G \times G' \rightarrow G$ supplies a $\mathfrak{D}N_p^K(Q)$ -interior algebra embedding

$$e_{\rho, Q^{\lambda}}^{K^{\lambda}}(B_{\ddot{E}, \varepsilon'}): \text{Ind}_{\rho_{Q^{\lambda}}}^{K^{\lambda}}(\bar{N}_{B_{\ddot{E}, \varepsilon'}}^{K^{\lambda}}(Q^{\lambda})) \rightarrow \bar{N}_{\text{Ind}_{\rho}(B_{\ddot{E}, \varepsilon'})}^K(Q).$$

Now, we have two idempotents in $(\text{Ind}_{\rho}(B_{\ddot{E}, \varepsilon'})(Q))^{\bar{N}_p^K(Q)}$, namely the two images $(\bar{N}_{\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}}^K(Q))(1)$ and $(e_{\rho, Q^{\lambda}}^{K^{\lambda}}(B_{\ddot{E}, \varepsilon'}))(1)$ of the respective unity elements of $\bar{N}_{\hat{A}_{\ddot{E}}}^K(Q)$ and $\text{Ind}_{\rho_{Q^{\lambda}}}^{K^{\lambda}}(\bar{N}_{B_{\ddot{E}, \varepsilon'}}^{K^{\lambda}}(Q^{\lambda}))$; moreover, since $(R_{\ddot{E}}, \ddot{R}_{\ddot{E}}, R'_{\varepsilon'})$ is basic, R stabilizes a basis in $\hat{A}_{\ddot{E}}$ [6, 17.1] and therefore the first idempotent is primitive. Consequently, $\bar{N}_{\tilde{h}_{\ddot{E}}^{\ddot{E}, \varepsilon'}}^K(Q)$ factorizes throughout $\bar{e}_{\rho, Q^{\lambda}}^{K^{\lambda}}(B_{\ddot{E}, \varepsilon'})$ determining a unique $\mathfrak{D}N_p^K(Q)$ -interior algebra exoembedding

$$\tilde{h}_{\ddot{E}}^{\lambda}(Q): \bar{N}_{\hat{A}_{\ddot{E}}}^K(Q) \rightarrow \text{Ind}_{\rho_{Q^{\lambda}}}^{K^{\lambda}}(\bar{N}_{B_{\ddot{E}, \varepsilon'}}^{K^{\lambda}}(Q^{\lambda})) \quad (4.7.1)$$

always compatible with the K -grading.

4.8. Once again because $(R_{\ddot{E}}, \ddot{R}_{\ddot{E}}, R'_{\varepsilon'})$ is basic, \ddot{R} stabilizes a basis in $\ddot{A}_{\ddot{E}}$ [6, 17.1] and, since $\ddot{A}_{\ddot{E}}$ is actually a matrix algebra over k (cf. 2.7), we claim that we can apply Proposition 3.7 to $\ddot{A}_{\ddot{E}}$ and $\text{Res}_{\rho'}((kG')_{\varepsilon'})$ obtaining a $\mathfrak{D}N_R^{K^{\lambda}}(Q^{\lambda})$ -interior algebra isomorphism

$$\bar{N}_{B_{\ddot{E}, \varepsilon'}}^{K^{\lambda}}(Q^{\lambda}) \cong \ddot{A}_{\ddot{E}}(Q^{\lambda}) \otimes_k \bar{N}_{\text{Res}_{\rho'}((kG')_{\varepsilon'})}^{K^{\lambda'}}(Q^{\lambda'})$$

where $K^{\lambda'}$ denotes the image of K in $\text{Aut}(Q^{\lambda'})$. Indeed, by the so-called *Frattini argument*, we have

$$Q \cdot N_G^K(Q) = Q \cdot C_G(Q) \cdot (N_G^K(Q) \cap N_G(T, h))$$

and therefore, since (T, h) is selfcentralizing, any $\varphi \in K$ is induced by some element $x \in N_{N_G^K(Q)}^K(T_v)$; then, it follows from Theorem 16.9 and Corollary 16.16 in [6] that there is $x' \in G'$ such that (x, x') normalizes the triple $(T_v, \ddot{T}_v, T'_{v'})$; thus, denoting by μ the restriction of λ to T , it is clear that

$\mu^{(x,x')}$ is also a section of the triple $(T_{\tilde{v}}, \tilde{T}_{\tilde{v}}, T'_{\nu'})$ and it follows from 2.12.1 that there is $y' \in G'$ fulfilling

$$\tilde{T}^{(1,y')} = \tilde{T} \quad \text{and} \quad \mu^{(x,x'y')} = \mu$$

and therefore $(x, x'y')$ normalizes T^λ and Q^λ and induces φ on Q^λ .

4.9. But, since $\tilde{T}_{\tilde{v}} \subset \tilde{R}_{\tilde{e}}$, we have a $\mathfrak{D}\tilde{T}$ -embedding $\tilde{A}_{\tilde{v}} \rightarrow \text{Res}_{\tilde{Q}}^{\tilde{R}}(\tilde{A}_{\tilde{e}})$ and therefore, since $\tilde{A}_{\tilde{v}}(T^\lambda) \neq \{0\}$ [6, 17.4.7] and $\tilde{A}_{\tilde{e}}$ admits an \tilde{R} -stable basis, we still have $\tilde{A}_{\tilde{e}}(Q^\lambda) \neq \{0\}$ [6, Lemma 7.10]; on the other hand, if \tilde{X} is a \tilde{T} -stable basis of $\tilde{A}_{\tilde{v}}$, it is easily checked that $(1 \times G') \otimes \tilde{X} \otimes (1 \times G')$ is a \tilde{T} -stable basis of $\text{Ind}_T^{T \times G'}(\tilde{A}_{\tilde{v}})$ and, in particular, it follows again from Corollary 5.8 in [4] that Q^λ has a unique local point χ^λ over the $k(T \times G')$ -interior algebra $\text{Ind}_T^{T \times G'}(\tilde{A}_{\tilde{v}})$, which necessarily comes from $\tilde{A}_{\tilde{v}}$; moreover, it is clear that $(1, y')$ acts on $\text{Ind}_T^{T \times G'}(\tilde{A}_{\tilde{v}})$ by conjugation, whereas the action of (x, x') on the group $T \times G'$ normalizing $\tilde{T}_{\tilde{v}}$ still determines a twisted action over this $k(T \times G')$ -interior algebra; at this point, the uniqueness of χ implies that the action of $(x, x'y')$ normalizes $(Q^\lambda)_{\chi^\lambda}$; that is to say, we get

$$\text{Res}_\varphi((\tilde{A}_{\tilde{e}})_{\chi^\lambda}) = \text{Res}_\varphi((\tilde{A}_{\tilde{v}})_{\chi^\lambda}) \cong (\tilde{A}_{\tilde{v}})_{\chi^\lambda} = (\tilde{A}_{\tilde{e}})_{\chi^\lambda}$$

which proves our claim.

4.10. Thus, setting $\tilde{D} = \tilde{A}_{\tilde{e}}(Q^\lambda)$ and $D' = \tilde{N}_{(kG')_{e'}}^{K\lambda'}(Q^{\lambda'})$, the $\mathfrak{D}N_P^K(Q)$ -interior algebra exoembedding (4.7.1) becomes

$$\tilde{h}_{\tilde{e}}^{\lambda'}(Q) : \tilde{N}_{\tilde{A}_{\tilde{e}}}^K(Q) \rightarrow \text{Ind}_{\rho_{Q^\lambda}^{K\lambda}}(\tilde{D} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D')); \quad (4.10.1)$$

once again, since $(\tilde{h}_{\tilde{e}}^{\lambda'}(Q))(1)$ belongs to a local point of $N_P^K(Q)$, it is clear that, for suitable points $\tilde{\beta}$ of $N_{\tilde{R}}^{K\lambda}(Q^\lambda)$ over \tilde{D} and β' of $N_{p'}^{K\lambda'}(Q^{\lambda'})$ over D' , $\tilde{h}_{\tilde{e}}^{\lambda'}(Q)$ factorizes throughout the canonical exoembedding

$$\text{Ind}_{\rho_{Q^\lambda}^{K\lambda}}(\tilde{D}_{\tilde{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'_{\beta'})) \rightarrow \text{Ind}_{\rho_{Q^\lambda}^{K\lambda}}(\tilde{D} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'))$$

determining a unique $\mathfrak{D}N_P^K(Q)$ -interior algebra exoembedding

$$\tilde{h}_{\tilde{e}}^{\tilde{\beta}, \beta'}(Q) : \tilde{N}_{\tilde{A}_{\tilde{e}}}^K(Q) \rightarrow \text{Ind}_{\rho_{Q^\lambda}^{K\lambda}}(\tilde{D}_{\tilde{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'_{\beta'})). \quad (4.10.2)$$

4.11. We claim that β' is a local point; indeed, let $T'_{\eta'}$ be a defect pointed group of $N_{p'}^{K\lambda'}(Q^{\lambda'})_{\beta'}$; then, it follows from Theorem 2.4 that there exists a $\mathfrak{D}N_{p'}^{K\lambda'}(Q^{\lambda'})$ -interior algebra exoembedding

$$\tilde{h}_{\beta'}^{\eta'} : D'_{\beta'} \rightarrow \text{Ind}_{T'_{\eta'}}^{N_{p'}^{K\lambda'}(Q^{\lambda'})}(D'_{\eta'});$$

note that, since $N_{p'}^{K\lambda'}(Q^{\lambda'})$ is a p -group, $\tilde{h}_{\beta'}^{\eta'}$ is actually an exoisomorphism [6, 2.12.2]. Consequently, setting $\tilde{T} = \rho'^{-1}(N_{p'}^{K\lambda'}(Q^{\lambda'}))$ and denoting by $\tau' : \tilde{T} \rightarrow T'$ the homomorphism determined by ρ'

(cf. 4.6), it follows from Proposition 12.12 in [6] that we have

$$\text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'_{\beta'}) \cong \text{Ind}_{\tilde{T}}^{N_{\tilde{R}}^{K\lambda}(Q^{\lambda})}(\text{Res}_{\tau'}(D'_{\eta'}));$$

then, by Corollary 12.7 and Proposition 12.9 in [6], we get a $\mathfrak{D}N_P^K(Q)$ -interior algebra exoembedding

$$\tilde{h}_{\tilde{\varepsilon}}^{\tilde{\eta}, \beta'}(Q) : \bar{N}_{\tilde{A}_{\tilde{\varepsilon}}}^K(Q) \rightarrow \text{Ind}_{\tilde{T}}^{N_{\tilde{R}}^{K\lambda}(Q^{\lambda})}(\ddot{D}_{\tilde{\beta}}) \otimes_k \text{Res}_{\tau'}(D'_{\eta'})$$

where we denote by $\tau : \tilde{T} \rightarrow N_P^K(Q)$ the group homomorphism determined by ρ (cf. 4.6).

4.12. Since the unity element forms a local point of $N_P^K(Q)$ on $\bar{N}_{\tilde{A}_{\tilde{\varepsilon}}}^K(Q)$, we necessarily have $\tau(\tilde{T}) = N_P^K(Q)$ and therefore we still have

$$N_{\tilde{R}}^{K\lambda}(Q^{\lambda}) = N_{\text{Ker}(\rho)}^{K\lambda}(Q^{\lambda}) \cdot \tilde{T};$$

moreover, since $\tilde{T} \subset N_{\tilde{R}}^{K\lambda}(Q^{\lambda})$ stabilizes a basis of $\ddot{D}_{\tilde{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'_{\beta'})$ and $\ddot{D}_{\tilde{\beta}}$ is projective as $\text{Ker}(\tau)$ -module, it follows from Lemma 13.6 in [6] that $\text{Ker}(\tau)$ has at least one complement \tilde{U} in \tilde{T} such that

$$(\ddot{D}_{\tilde{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(D'_{\beta'}))(\tilde{U}) \neq \{0\};$$

thus, since $\ddot{D}_{\tilde{\beta}}$ is also projective as $\text{Ker}(\tau')$ -module, we have $\tilde{U} \cap \text{Ker}(\tau') = \{1\}$ and therefore \tilde{U} maps into T' , so that $|N_P^K(Q)| = |\tilde{U}| \leq |T'|$ which forces the equalities $T' = N_{P'}^{K\lambda'}(Q^{\lambda'})$ and $\eta' = \beta'$.

4.13. Furthermore, since $(Q^{\lambda'}, f^{\lambda'}) \subset (R', g')$, we have [6, 2.13.2]

$$f^{\lambda'} \text{Br}_Q(\varepsilon') = \text{Br}_Q(\varepsilon');$$

hence, $N_{P'}^{K\lambda'}(Q^{\lambda'})_{\beta'}$ is a maximal local pointed group over $kN_{G'}^{K\lambda'}(Q^{\lambda'})f^{\lambda'}$ (cf. 4.5), so that $(kN_{G'}^{K\lambda'}(Q^{\lambda'}))_{\beta'}$ is a source algebra of $kN_{G'}^{K\lambda'}(Q^{\lambda'})f^{\lambda'}$, and the $\mathfrak{D}N_P^K(Q)$ -interior algebra exoembedding (4.10.2) becomes

$$\tilde{h}_{\tilde{\varepsilon}}^{\tilde{\beta}, \beta'}(Q) : \bar{N}_{\tilde{A}_{\tilde{\varepsilon}}}^K(Q) \rightarrow \text{Ind}_{\rho_{Q^{\lambda}}^{K\lambda}}(\ddot{D}_{\tilde{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(kN_{G'}^{K\lambda'}(Q^{\lambda'}))_{\beta'}). \quad (4.13.1)$$

4.14. On the other hand, it follows from Lemma 4.18 below that there is a local point $\bar{\varepsilon}$ of $N_R^K(Q)$ over $kN_G^K(Q)f$ such that the structural homomorphism from $N_G^K(Q)$ to $N_{kG}^K(Q)$ induces a $kN_R^K(Q)$ -interior algebra embedding

$$(kN_G^K(Q))_{\bar{\varepsilon}} \rightarrow \bar{N}_{(kG)_{\bar{\varepsilon}}}^K(Q); \quad (4.14.1)$$

moreover, according to our choice of R_{ε} (cf. 4.2) and to the same lemma, $N_R^K(Q)_{\bar{\varepsilon}}$ is a maximal local pointed group over $kN_G^K(Q)f$ and therefore $N_P^K(Q)$ is a defect group of f as a block of $N_G^K(Q)$, and the $kN_P^K(Q)$ -interior algebra $(kN_G^K(Q))_{\bar{\varepsilon}}$ is a source algebra of $kN_G^K(Q)f$.

4.15. Finally, according to our hypothesis (cf. 2.12, 4.1 and 4.3), $\hat{A}_{\hat{\varepsilon}}$ is a 0-split $\mathfrak{D}(P \times P)$ -module and it follows from Proposition 2.8 that we have a kR -interior algebra isomorphism

$$(kG)_{\varepsilon} \cong H_0(\hat{A}_{\hat{\varepsilon}});$$

then, by Lemma 3.11 applied to the kR -interior algebra $\hat{A}_{\hat{\varepsilon}}$, $\bar{N}_{\hat{A}_{\hat{\varepsilon}}}^K(Q)$ is a 0-split $\mathfrak{D}(N_P^K(Q) \times N_P^K(Q))$ -module and we get a $kN_P^K(Q)$ -interior algebra embedding (cf. (4.14.1))

$$(kN_G^K(Q))_{\bar{\varepsilon}} \rightarrow \bar{N}_{(kG)_{\varepsilon}}^K(Q) \cong H_0(\bar{N}_{\hat{A}_{\hat{\varepsilon}}}^K(Q)).$$

4.16. That is to say, denoting by $\hat{\varepsilon}$ the local point of $N_P^K(Q)$ over

$$\text{Ind}_{\rho_{Q^{\lambda}}^{K\lambda}}(\ddot{D}_{\ddot{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(kN_{G'}^{K\lambda'}(Q^{\lambda'}))_{\beta'})$$

determined by the local point $\bar{\varepsilon}$ and the embedding $\tilde{h}_{\bar{\varepsilon}}^{\ddot{\beta}, \beta'}(Q)$ (cf. Proposition 2.8 and (4.13.1)), the $kN_P^K(Q)$ -interior algebra

$$\text{Ind}_{\rho_{Q^{\lambda}}^{K\lambda}}(\ddot{D}_{\ddot{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(kN_{G'}^{K\lambda'}(Q^{\lambda'}))_{\beta'})_{\hat{\varepsilon}}$$

is a 0-split $\mathfrak{D}(N_P^K(Q) \times N_P^K(Q))$ -module and we have a $kN_P^K(Q)$ -interior algebra isomorphism

$$(kN_G^K(Q))_{\bar{\varepsilon}} \cong \mathbb{H}_0(\text{Ind}_{\rho_{Q^{\lambda}}^{K\lambda}}(\ddot{D}_{\ddot{\beta}} \otimes_k \text{Res}_{\rho_{Q^{\lambda'}}^{K\lambda'}}(kN_{G'}^{K\lambda'}(Q^{\lambda'}))_{\beta'})_{\hat{\varepsilon}}).$$

Consequently, it follows from Theorem 18.8 in [6] that the block algebras $kN_G^K(Q)f$ and $kN_{G'}^{K\lambda'}(Q^{\lambda'})f^{\lambda'}$ are Rickard equivalent.

4.17. Moreover, from our basic hypothesis, we know that $\ddot{A}_{\bar{\varepsilon}}$ admits $\ddot{R} \times_R \ddot{R}$ - and $\ddot{R} \times_{R'} \ddot{R}$ -stable bases \ddot{Y} and \ddot{Y}' where $\text{Ker}(\rho) \times \text{Ker}(\rho)$ and $\text{Ker}(\rho') \times \text{Ker}(\rho')$ respectively act freely; then, it is elementary to check that $\text{Br}_{Q^{\lambda}}(\ddot{Y}^{Q^{\lambda}})$ and $\text{Br}_{Q^{\lambda}}(\ddot{Y}'^{Q^{\lambda}})$ are bases of $\ddot{D} = \ddot{A}_{\bar{\varepsilon}}(Q^{\lambda})$, and, since we have

$$N_{\ddot{R}}^{K\lambda}(Q^{\lambda}) = C_{\ddot{R}}(Q^{\lambda}) \cdot \Delta_{\lambda}(N_R^K(Q)),$$

it is not difficult to check that $N_{\ddot{R}}^{K\lambda}(Q^{\lambda}) \times_{N_R^K(Q)} N_{\ddot{R}}^{K\lambda}(Q^{\lambda})$ stabilizes and $N_{\text{Ker}(\rho)}^{K\lambda}(Q^{\lambda}) \times_{N_{\text{Ker}(\rho)}^{K\lambda}(Q^{\lambda})} N_{\text{Ker}(\rho)}^{K\lambda}(Q^{\lambda})$ acts freely on the basis $\text{Br}_{Q^{\lambda}}(\ddot{Y}^{Q^{\lambda}})$; similarly, $N_{\ddot{R}}^{K\lambda}(Q^{\lambda}) \times_{N_{R^{\lambda'}}^{K\lambda'}(Q^{\lambda'})} N_{\ddot{R}}^{K\lambda}(Q^{\lambda})$ stabilizes and $N_{\text{Ker}(\rho')}^{K\lambda}(Q^{\lambda}) \times_{N_{\text{Ker}(\rho')}^{K\lambda}(Q^{\lambda})} N_{\text{Ker}(\rho')}^{K\lambda}(Q^{\lambda})$ acts freely on the basis $\text{Br}_{Q^{\lambda}}(\ddot{Y}'^{Q^{\lambda}})$. In conclusion, the block algebras $kN_G^K(Q)f$ and $kN_{G'}^{K\lambda'}(Q^{\lambda'})f^{\lambda'}$ are basic Rickard equivalent.

Lemma 4.18. Let (Q, f) be a Brauer (b, G) -pair, K a subgroup of the image of $N_G(Q, f)$ in $\text{Aut}(Q)$ and v the point of $Q \cdot N_G^K(Q)$ over kGb such that $\text{Br}_Q(v) = \{f\}$. For any local pointed group R_{ε} over kGb such that $Q \subset R$ and $R_{\varepsilon} \subset Q \cdot N_G^K(Q)_v$, there is a local point $\bar{\varepsilon}$ of $N_R^K(Q)$ over $kN_G^K(Q)f$ such that the structural homomorphism from $N_G^K(Q)$ to $N_R^K(Q)$ induces a $kN_R^K(Q)$ -interior algebra embedding

$$(kN_G^K(Q))_{\bar{\varepsilon}} \rightarrow \bar{N}_{(kG)_{\varepsilon}}^K(Q). \quad (4.18.1)$$

Moreover, $N_R^K(Q)_{\bar{\varepsilon}}$ is a maximal local pointed group over $kN_G^K(Q)f$ if and only if R_{ε} is a defect pointed group of $Q \cdot N_G^K(Q)_{\nu}$. In particular, $N_R^K(Q)$ is a defect group of f as a block of $N_G^K(Q)$ if and only if R is a defect group of f as a block of $Q \cdot N_G^K(Q)$.

Proof. As usual, we identify $(kG)(Q)$ with $kC_G(Q)$; from the existence of stable bases, it is easily checked that $\text{Br}((kG)^H) = kC_G(Q)^H$ for any subgroup H of $N_G(Q)$ containing Q ; hence, for any point β of H over kG , if $\text{Br}_Q(\beta) \neq \{0\}$ then $\text{Br}_Q(\beta)$ is a point of H over $kC_G(Q)$ and, conversely, any point of H over $kC_G(Q)$ can be lifted to $(kG)^H$; note that, since $Q \cdot N_G^K(Q)$ contains $C_G(Q)$, $kC_G(Q)^{Q \cdot N_G^K(Q)}$ is contained in $Z(kC_G(Q))$, which guarantees the existence of ν . Thus, since $\text{Br}_R(\varepsilon) \neq \{0\}$ and $R = Q \cdot N_R^K(Q)$, $\text{Br}_Q(\varepsilon)$ is a local point of both R and $N_R^K(Q)$ over $kC_G(Q)$, and the inclusion of R_{ε} in $Q \cdot N_G^K(Q)_{\nu}$ forces

$$f \text{Br}_Q(\varepsilon) = \text{Br}_Q(\varepsilon).$$

In particular, choosing $i \in \varepsilon$ such that $(kG)_{\varepsilon} = i(kG)i$, $\text{Br}_Q(i)$ is a primitive idempotent in $(kC_G(Q)f)^{N_R^K(Q)}$ and, since the inclusion of $C_G(Q)$ in $N_G^K(Q)$ induces an injective homomorphism

$$(kC_G(Q)f)(N_R^K(Q)) \rightarrow (kN_G^K(Q)f)(N_R^K(Q)), \quad (4.18.2)$$

there is a primitive idempotent j in $(kN_G^K(Q)f)^{N_R^K(Q)}$ fulfilling

$$j \text{Br}_Q(i) = j = \text{Br}_Q(i)j \quad \text{and} \quad \text{Br}_{N_R^K(Q)}^{kN_G^K(Q)f}(j) \neq 0.$$

That is to say, j belongs to a local point $\bar{\varepsilon}$ of $N_R^K(Q)$; moreover, it is clear that the $kN_G^K(Q)$ -interior algebra isomorphism $kN_G^K(Q) \cong \tilde{N}_{kG}^K(Q)$ [7, Proposition 3.5] maps $(kN_G^K(Q))_{\bar{\varepsilon}}$ bijectively onto $j\tilde{N}_{kG}^K(Q)j$, whereas we have

$$N_{(kG)_{\varepsilon}}^K(Q) = \bigoplus_{\varphi \in K} (i(kG)i)(\Delta_{\varphi}(Q)) = \text{Br}_Q(i)\tilde{N}_{kG}^K(Q)\text{Br}_Q(i)$$

and embedding (4.18.1) easily follows.

Conversely, for any local point $\bar{\varepsilon}'$ of $N_R^K(Q)$ over $kN_G^K(Q)f$, it is clear that there are $j' \in \bar{\varepsilon}'$ and a primitive idempotent \bar{i}' in

$$(kC_G(Q)f)^{N_R^K(Q)} = (kC_G(Q)f)^R$$

fulfilling $j'\bar{i}' = j' = \bar{i}'j'$, and therefore we have $\bar{i}' = \text{Br}_Q(i')$ for a suitable primitive idempotent in $(kGb)^R$ belonging to a local point ε' of R over kGb such that $R_{\varepsilon'} \subset Q \cdot N_G^K(Q)_{\nu}$. Hence, since we have $R = Q \cdot N_R^K(Q)$, so that R and $N_R^K(Q)$ determine each other, and all the maximal local pointed groups contained in a pointed group are mutually conjugate, it is easily checked that $N_R^K(Q)_{\bar{\varepsilon}}$ is a maximal local pointed group over $kN_G^K(Q_{\delta})f$ if and only if R_{ε} is a maximal local pointed group over $Q \cdot N_G^K(Q_{\delta})_{\nu}$. The last statement follows from this one applied once again to $K \cdot \text{Inn}(Q)$. \square

References

- [1] A. Marcus, On equivalences between blocks of group algebras: Reduction to the simple components, J. Algebra 184 (1996) 372–396.
- [2] L. Puig, Pointed groups and construction of characters, Math. Z. 176 (1981) 265–292.

- [3] L. Puig, Local extensions in endo-permutation modules split: A proof of Dade's theorem, in: *Seminaire sur les Groupes Finis III*, Publ. Math. Paris VII 25 (1986).
- [4] L. Puig, Nilpotent blocks and their source algebras, *Invent. Math.* 93 (1988) 77–116.
- [5] L. Puig, Affirmative answer to a question of Feit, *J. Algebra* 131 (1990) 513–526.
- [6] L. Puig, On the local structure of Morita and Rickard equivalences between Brauer blocks, *Progr. Math.* 178 (1999).
- [7] L. Puig, Y. Zhou, A local property of basic Morita equivalences, *Math. Z.* 256 (2007) 551–562.
- [8] J. Rickard, Morita theory for derived categories, *J. Lond. Math. Soc.* 39 (1989) 436–456.
- [9] J. Rickard, Splendid equivalences: Derived categories and permutation modules, *Proc. Lond. Math. Soc.* 72 (1996) 331–358.